



Glyphosate

herbicide



“..... A Growing Partnership With Nature”

TABLE OF CONTENTS

Precautionary Statements	1	Cropping Systems	21
General Information	2	Row Crops, Cereal Grains, Citrus, Tree Nuts, Vine Crops, Tree Fruits, Vegetables, Small Fruits and Berries, Forage Crops, and Legumes, Tropical Crops, Sapote	22
Information on Resistant Weeds	3	Asparagus	23
Application Information	3	Berries and Small Fruits	24
Mixing, Additives and Application Instructions	3	Corn	24
Application Equipment & Techniques	4	Fallow and Reduced Tillage Systems	24
Aerial Equipment	4	Tank Mixtures	24
Broadcast Equipment	4	Ecofarming Systems	25
Controlled Droplet Application (CDA)	4	Aid to Tillage	25
Hand-Held and High Volume Equipment	5	Postharvest Grain Sorghum, Sorghum Regrowth Control	26
Selective Equipment	5	Pastures	26
Shielded Sprayers	5	Sugarcane	26
Wiper Applications	5	Conservation Tillage, Minimum Tillage and No-Till Systems	26
Low-Volume Broadcast Application (Low Rate Technology)	6	Corn and Soybeans Tank Mixtures	27
Tank Mixtures	8	Preharvest Applications	27
High-Volume Broadcast Applications	8	Soybeans	27
Perennial Weeds	8	Cotton	27
Woody Brush and Trees	12	Grain Sorghum	27
Noncrop Uses	14	Wheat	28
Industrial Recreational and Public Areas	14	Tree and Vine Crops	28
Tank Mixtures for Industrial Sites & Forestry Site Preparation	14	Middle Management for Annual Weeds in Middles	28
Tank Mixtures for Noncrop Sites	15	Between Rows of Trees and Vine Crops Strips for Annual and Perennial Weeds in Strips of Tree and Vine Crops Tank Mixtures with Residual Herbicides	29
Farmstead Weed Control	15	Perennial Grass Suppression Orchard Floors	29
Farm Ditches	15	Low Volume Application (Florida and Texas)	30
Conservation Reserve Program (CRP Acres)	15	Tree Crops	30
Dormant Rangeland	16	Vine Crops	31
Habitat Management	16	For Cotton with the "Roundup Ready" Gene in Crop Applications	31
Christmas Trees	16	Tank Mixtures	32
Silvicultural Sites and Rights-of-Way	16	GLYPHOSATE Plus STAPLE®	32
Site Preparation	17	Sprayer Preparation	33
Postdirected Spray	17	Sprayer Cleanup	33
Conifer Release	17	Spray Drift Management	34
Cut Stump Treatments	18	Importance of Droplet Size	34
Injection and Frill Applications	18	Controlling Droplet Size	34
Turfgrasses and Grasses for Seed Production	18	General Techniques	34
Preplant and Renovation	18	Aircraft	34
Turfgrasses	18	Boom Length and Height	34
Grasses for Seed Production	19	Wind	34
Annual Weed Control in Dormant Bermudagrass and Bahia Grass Turf	19	Temperature and Humidity	34
Release of Bermudagrass or Bahia Grass	19	Surface Temperature Inversions	34
Release of Actively Growing Bermudagrass	20	Shielded Sprayers	34
Cool Season Turf Growth Regulation	20	For Aerial Application in Arkansas Only	34
Annual Grasses	21	For Aerial Application in California Only	35
Tank Mixtures	21	For Aerial Application in Fresno County, CA Only	35
Tall Fescue	21	Storage and Disposal	36
Smooth Brome	21	Limitation of Warranty and Liability	37
Bahia Grass Seedhead and Vegetative Suppression	21		



Glyphosate

herbicide

Active IngredientBy Weight

*Glyphosate, N-(phosphonomethyl) 41%
Glycine, in the form of its isopropylamine salt

Inert Ingredients 59%

TOTAL100%

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

EPA Reg. No. 352-607

Net Contents: 2.5 gal./30 gal.

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Immediately hold eyelids open and flush with plenty of water for at least 15 minutes. Get medical attention.

IF SWALLOWED: This product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

IF INHALED: Remove individual to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

For medical emergencies involving this product, call toll free 1-800-441-3637.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

WARNING! CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY. HARMFUL IF SWALLOWED OR INHALED. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.
- Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL/CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.
- Protective eyewear.

Do not apply this product through any type of irrigation system.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

GENERAL INFORMATION

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

This product, a water soluble liquid, mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise specified on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "WEEDS CONTROLLED" section of this label. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per acre within the recommended range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (noncultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

INFORMATION ON RESISTANT WEEDS

When herbicides with the same mode of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant weed biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. These resistant weed biotypes may not be adequately controlled. Cultural practices such as tillage, preventing weed escapes from going to seed, and using herbicides with different modes of action within and between crop seasons can aid in delaying the proliferation and possible dominance of herbicide resistant weed biotypes.

APPLICATION INFORMATION

MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. DO NOT APPLY WHEN WIND OR OTHER CONDITIONS FAVOR DRIFT. HAND-GUN APPLICATIONS SHOULD BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS. NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

MIXING

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the recommended amount of this product (see the "DIRECTIONS FOR USE" and "WEEDS CONTROLLED" sections of this label) near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

TANK MIXTURES

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Mix labeled tank mixtures of this product with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it SLOWLY through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
6. Continue filling the spray tank with water and add the required amount of this product near the end of the filling process.
7. Where nonionic surfactant is recommended, add this to the spray tank before completing the filling process.
8. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water soluble liquid followed by surfactant.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles.

Clean sprayer and parts immediately after using this product by thoroughly flushing with water.

ADDITIVES

SURFACTANTS: Nonionic surfactants which are labeled for use with herbicides may be used. Do not reduce rates of this product when adding surfactant. When adding additional surfactant, use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

AMMONIUM SULFATE: The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product, and this product plus 2,4-D, "Banvel" or residual herbicide tank mixtures on annual and perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Low-quality ammonium sulfate may contain material that will not readily dissolve, which could result in nozzle tip plugging. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet line. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides or surfactant. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: The use of ammonium sulfate as an additive does not preclude the need for additional surfactant. Do not use herbicide rates lower than recommended in this label.

COLORANTS OR DYES

Agriculturally-approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's recommendations.

APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

Aerial* - Fixed Wing and Helicopter (for aerial applications in the states of Arkansas and California see additional instructions at end of the label).

Broadcast Spray

Controlled Droplet Applicator (CDA) - Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

Hand-Held and High-Volume Spray Equipment - Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers**, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

Selective Equipment - Recirculating sprayers, shielded sprayers and wiper applicators.

PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

* This product plus DuPont OUST®, "Banvel" or 2,4-D tank mixtures may not be applied by air in California.

** This product is not registered in California or Arizona for use in mistblowers.

AERIAL EQUIPMENT

Use the recommended rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. See the "WEEDS CONTROLLED" section of this label for specific rates. Unless otherwise specified, do not exceed 1 quart per acre. Aerial applications of this product may be made in annual cropping conventional tillage systems, fallow and reduced tillage systems and preharvest applications. Refer to the individual use area sections of this label for recommended volumes and application rates. **FOR AERIAL APPLICATION IN CALIFORNIA OR ARKANSAS, REFER TO THE ADDITIONAL INSTRUCTIONS AT THE END OF THIS LABEL.**

BROADCAST EQUIPMENT

For control of annual or perennial weeds listed on this label using broadcast equipment -- Use the recommended rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label. See the "WEEDS CONTROLLED" section of this label for specific rates. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

CONTROLLED DROPLET APPLICATION (CDA)

The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre.

For the control of labeled annual weeds with hand-held CDA units, apply a 20 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 MPH (1 quart per acre). For the control of labeled perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2 to 4 quarts per acre).

Controlled droplet application equipment produces a spray pattern which is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

HAND-HELD AND HIGH-VOLUME EQUIPMENT

USE COARSE SPRAYS ONLY

Mix this product in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff.

For control of annual weeds listed on this label, apply a 0.5 percent solution of this product plus nonionic surfactant to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. Allow three or more days before tillage or mowing.

For annual weeds over 6 inches tall, or when not using additional surfactant, or unless otherwise specified, use a 1 percent solution. For best results, use a 2 percent solution on harder-to-control perennials, such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

When using application methods which result in less than complete coverage, use a 5 percent solution for annual and perennial weeds and a 5 to 10 percent solution for woody brush and trees.

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

SPRAY SOLUTION

AMOUNT OF GLYPHOSATE

DESIRED VOLUME	1/2%	1%	1 1/2%	2%	5%	10%
1 GAL	2/3 OZ	1-1/3 OZ	2 OZ	2-2/3 OZ	6-1/2 OZ	13 OZ
25 GAL	1 PT	1 QT	1-1/2 QT	2 QT	5 QT	10 QT
100 GAL	2 QT	1 GAL	1-1/2 GAL	2 GAL	5 GAL	10 GAL

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the recommended amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

SELECTIVE EQUIPMENT

This product may be applied through a recirculating spray system, a shielded applicator, or a wiper applicator after dilution and thorough mixing with water to listed weeds growing in any noncrop site specified on this label and only when specifically recommended in cropping systems.

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

A shielded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide.

A wiper applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution.

AVOID CONTACT WITH DESIRABLE VEGETATION.

Contact of the herbicide solution with the desirable vegetation may result in damage or destruction. Applicators used above desired vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam, or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

SHIELDED APPLICATORS

When applied as directed under conditions described for shielded applicators, this product will control those weeds listed in the "WEEDS CONTROLLED" section of this label.

Use the following equation to convert from a broadcast rate per acre to a band rate per acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE per acre} = \text{Herbicide Band RATE per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME of solution per acre} = \text{Band VOLUME of solution per acre}$$

Use nozzles that provide uniform coverage within the treated area. Keep shields on shielded sprayers adjusted to protect desirable vegetation. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT WITH DESIRABLE VEGETATION.**

For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

WIPER APPLICATORS

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

For Rope or Sponge Wick Applicators: Mix 1 gallon of this product in 2 gallons of water to prepare a 33 percent solution. Apply this solution to weeds listed in this "WIPER APPLICATORS" section.

For Porous-Plastic Applicators: Solutions ranging from 33 to 100 percent of this product in water may be used in porous-plastic wiper applicators.

When applied as recommended under the conditions described for "WIPER APPLICATORS", this product CONTROLS the following weeds:

ANNUAL GRASSES

Corn, *Zea mays*

Panicum, Texas, *Panicum texanum*

Rye, common, *Secale cereale*

Shattercane, *Sorghum bicolor*

ANNUAL BROADLEAVES

Sicklepod, *Cassia obtusifolia*

Spanishneedles, *Bidens bipinnata*

Starbur, bristly, *Acanthospermum hispidum*

When applied as recommended under the conditions described for "WIPER APPLICATORS", this product SUPPRESSES the following weeds:

ANNUAL BROADLEAVES

Beggarweed, Florida, *Desmodium tortuosum*

Dogfennel, *Eupatorium capilliflorum*

Pigweed, redroot, *Amaranthus retroflexus*

Ragweed, common, *Ambrosia artemisiifolia*

Ragweed, giant, *Ambrosia trifida*

Sunflower, *Helianthus annuus*

Thistle, musk, *Carduus nutans*

Velvetleaf, *Abutilon theophrasti*

PERENNIAL GRASSES

Bermudagrass, *Cynodon dactylon*

Guineagrass, *Panicum maximum*

Johnsongrass, *Sorghum halepense*

Smutgrass, *Sporobolus poiretii*

Vaseygrass, *Paspalum urvillei*

PERENNIAL BROADLEAVES

Dogbane, hemp, *Apocynum cannabinum*

Milkweed, *Asclepias syriaca*

Nightshade, silverleaf, *Solanum elaeagnifolium*

Thistle, Canada, *Cirsium arvense*

WEEDS CONTROLLED

This herbicide controls many annual and perennial grasses and broadleaf weeds.

ANNUAL WEEDS

- Apply to actively growing grass and broadleaf weeds.
- Allow at least 3 days after treatment before tillage.
- For maximum agronomic benefit, apply when weeds are 6 inches or less in height.
- To prevent seed production, applications should be made prior to seedhead formation.
- This product does not provide residual control; therefore, delay application until maximum weed emergence. Repeat treatments may be necessary to control later germinating weeds.

LOW-VOLUME BROADCAST APPLICATION (LOW-RATE TECHNOLOGY)

When applied as directed under the conditions described, this product will control the weeds listed below when:

1. Water carrier volumes of 3 to 10 gallons per acre for ground applications and 3 to 5 gallons per acre for aerial applications are recommended. (See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for approved sites.)
2. A nonionic surfactant is added at 0.5 to 1 percent by total spray volume. Use 0.5 percent surfactant concentration when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration for those surfactants containing less than 70 percent active ingredient.

NOTE: The addition of 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water may increase the performance of this product on annual weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label.

Do not tank-mix with soil residual herbicides when using these rates unless otherwise specified.

For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Refer to the "TANK MIXTURES" portion of this section for control of additional broadleaf weeds.

WEED SPECIES	MAXIMUM HEIGHT/LENGTH	RATE PER ACRE* (Fluid Ounces)
Foxtail, <i>Setaria spp</i>	12"	8 oz.
Barnyardgrass, <i>Echinochloa crus-galli</i>	6" (0 to 4" 4 to 6")	12 oz. 16 oz. ¹ 24 oz. ¹)
Bluegrass, annual, <i>Poa annua</i>		
Brome, downy** <i>Bromus tectorum</i>		
Mustard, blue <i>Chorispora tenella</i>		
Mustard, tansy <i>Descurainia pinnata</i>		
Mustard, tumble <i>Sisymbrium altissimum</i>		
Mustard, wild <i>Sinapis arvensis</i>		
Spurry, umbrella <i>Holosteum umbellatum</i>		
Barley <i>Hordeum vulgare</i>	12"	
Rye <i>Secale cereale</i>		
Sandbur, field <i>Cenchrus spp.</i>		
Shattercane <i>Sorghum bicolor</i>		
Stinkgrass <i>Eragrostis cilianensis</i>		
Wheat <i>Triticum aestivum</i>	18"	
Morningglory <i>Ipomoea spp</i>	2"	16 oz.
Sicklepod <i>Cassia obtusifolia</i>		
Bluegrass, bulbous <i>Poa bulbosa</i>	6"	
Cheat <i>Bromus secalinus</i>		
Chickweed, common <i>Stellaria media</i>		
Chickweed, mouseear <i>Cerastium vulgatum</i>		
Corn <i>Zea mays</i>		
Goatgrass, jointed <i>Aegilops cylindrica</i>		
Groundsel, common <i>Senecio vulgaris</i>		
Henbit <i>Lamium amplexicaule</i>		
Horseweed/Marestail <i>Conyza canadensis</i>		
Lambsquarters, common <i>Chenopodium album</i>		
Pennycress, field <i>Fanweed Thlaspi arvense</i>		
Rocket, London <i>Sisymbrium irio</i>		
Ryegrass, Italian <i>Lolium multiflorum</i>		
Shepherdspurse <i>Capsella bursa-pastoris</i>		
Spurge, annual <i>Euphorbia spp</i>		
Buttercup <i>Ranunculus spp</i>	12"	

WEED SPECIES	MAXIMUM HEIGHT/LENGTH	RATE PER ACRE* (Fluid Ounces)
Cocklebur <i>Xanthium strumarium</i>		
Crabgrass <i>Digitaria spp</i>		
Dwarf dandelion <i>Krigia cespitosa</i>		
Falseflax, smallseed <i>Camelina microcarpa</i>		
Foxtail, Carolina <i>Alopecurus carolinianus</i>		
Johnsongrass, seedling <i>Sorghum halepense</i>		
Oats, wild <i>Avena fatua</i>		
Panicum, fall <i>Panicum dichotomiflorum</i>		
Panicum, Texas <i>Panicum texanum</i>		
Pigweed, redroot <i>Amaranthus retroflexus</i>		
Pigweed, smooth <i>Amaranthus hybridus</i>		
Witchgrass <i>Panicum capillare</i>		
Sicklepod <i>Cassia obtusifolia</i>	3 to 4"	24 oz
Signalgrass, broadleaf <i>Brachiaria platyphylla</i>	4"	
Horseweed/Marestail <i>Conyza canadensis</i>	7 to 12"	
Lambsquarters, common <i>Chenopodium album</i>		
Spurge, annual <i>Euphorbia spp.</i>		
Rice, red <i>Oryza sativa</i>	4"	32 oz.
Teaweed <i>Sida spinosa</i>		
Sprangletop <i>Leptochloa spp</i>	6"	
Geranium, Carolina <i>Geranium carolinianum</i>	12"	
Goosegrass <i>Eleusine indica</i>		
Primrose, cutleaf evening <i>Oenothera laciniata</i>		
Pusley, Florida <i>Richardia scabra</i>		
Sicklepod <i>Cassia obtusifolia</i>	5 to 12"	
Spanishneedles <i>Bidens bipinnata</i>		
Filaree <i>Erodium spp</i>	12"	48 oz
Sprangletop <i>Leptochloa spp</i>		

¹ Use these rates to control barnyardgrass in Alabama, Arkansas, Mississippi, Missouri, Louisiana and Texas for preplant treatments.

* For those rates less than 32 fluid ounces per acre, this product at rates up to 32 fluid ounces per acre may be used where heavy weed densities exist.

** For control in no-till systems, use 16 fluid ounces per acre.

TANK MIXTURES

GLYPHOSATE plus "Banvel" plus Nonionic Surfactant

GLYPHOSATE plus 2,4-D plus Nonionic Surfactant

DO NOT APPLY "BANVEL" OR 2,4-D TANK MIXTURES BY AIR IN CALIFORNIA.

These tank mixtures are recommended for use in fallow and reduced tillage areas only. Follow use directions as given in the "LOW-VOLUME BROADCAST APPLICATION" section.

This product plus "Banvel" or 2,4-D will control the annual grasses and broadleaf weeds listed for this product alone at the indicated heights (except 8 fluid ounces per acre applications), plus the following broadleaf weeds. For those weeds previously listed at 8 fluid ounces of this product alone per acre, use 12 fluid ounces in these tank mixtures.

NOTE: Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if "Banvel" is applied within 45 days of planting. The addition of "Banvel" in a mixture with this product may provide short-term residual control of selected weed species.

Apply 12 to 16 fluid ounces of this product plus 0.25 pound a.i. of "Banvel" or 0.5 pound a.i. of 2,4-D, plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre to control dense populations of the following annual broadleaf weeds when less than the height indicated:

Cocklebur (12"),	<i>Xanthium strumarium</i>
Kochia* (6"),	<i>Kochia scoparia</i>
Lambsquarters (12"),	<i>Chenopodium album</i>
Lettuce, prickly (6"),	<i>Lactuca serriola</i>
Marestail/Horseweed (6"),	<i>Conyza canadensis</i>
Morningglory (6"),	<i>Ipomoea spp.</i>
Pigweed, redroot (12"),	<i>Amaranthus retroflexus</i>
Pigweed, smooth (12"),	<i>Amaranthus hybridus</i>
Thistle, Russian (12"),	<i>Salsola kali</i>

*Controlled with "Banvel" tank mixture only.

Apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D, plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre to control the following annual broadleaf weeds when less than 6 inches in height.

Ragweed, common,	<i>Ambrosia artemisiifolia</i>
Ragweed, giant,	<i>Ambrosia trifida</i>
Smartweed, Pennsylvania,	<i>Polygonum pensylvanicum</i>
Velvetleaf,	<i>Abutilon theophrasti</i>

HIGH-VOLUME BROADCAST APPLICATIONS

When applied as directed under the conditions described, this product will control the weeds listed below when water carrier volumes are 10 to 40 gallons per acre for ground applications.

Apply 1 to 1.5 quarts of this product per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume. Use 1 quart per acre if weeds are less than 6 inches tall and 1.5 quarts per acre if weeds are over 6 inches tall. If weeds have been mowed, grazed or cut, allow adequate time for new growth to reach recommended stages prior to treatment. These rates will also provide control of weeds listed in the "LOW-VOLUME BROADCAST APPLICATION" section.

WEED SPECIES

Balsamapple*,	<i>Momordica charantia</i>
Bassia, fivehook,	<i>Bassia hyssopifolia</i>
Brome,	<i>Bromus spp.</i>
Fiddleneck,	<i>Amsinckia spp.</i>
Fleabane, hairy,	<i>Conyza bonariensis</i>
Fleabane,	<i>Erigeron spp.</i>
Kochia,	<i>Kochia scoparia</i>
Lettuce, prickly,	<i>Lactuca serriola</i>
Panicum,	<i>Panicum spp.</i>
Ragweed, common,	<i>Ambrosia artemisiifolia</i>
Ragweed, giant,	<i>Ambrosia trifida</i>
Smartweed, Pennsylvania,	<i>Polygonum pensylvanicum</i>
Sowthistle, annual,	<i>Sonchus oleraceus</i>
Sunflower,	<i>Helianthus annuus</i>
Thistle, Russian,	<i>Salsola kali</i>
Velvetleaf,	<i>Abutilon theophrasti</i>

*Apply with hand-held equipment only.

PERENNIAL WEEDS

Apply this product as follows to control or destroy most perennial weeds:

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product on perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label.

When applied as recommended under the conditions described, this product WILL CONTROL the following PERENNIAL WEEDS:

Alfalfa,	<i>Medicago sativa</i>
Alligatorweed*,	<i>Alternanthera philoxeroides</i>
Anise (fennel),	<i>Foeniculum vulgare</i>
Artichoke, Jerusalem,	<i>Helianthus tuberosus</i>
Bahiagrass,	<i>Paspalum notatum</i>
Bentgrass,	<i>Agrostis spp.</i>
Bermudagrass,	<i>Cynodon dactylon</i>
Bermudagrass, water (knotgrass),	<i>Paspalum distichum</i>
Bindweed, field,	<i>Convolvulus arvensis</i>
Bluegrass, Kentucky,	<i>Poa spp.</i>
Blueweed, Texas,	<i>Helianthus ciliaris</i>
Brackenfern,	<i>Pteridium aquilinum</i>
Bromegrass, smooth,	<i>Bromus inermis</i>
Bursage, woollyleaf,	<i>Franseria tomentosa</i>
Canarygrass, reed,	<i>Phalaris arundinacea</i>
Cattail,	<i>Typha spp.</i>
Clover, red,	<i>Trifolium pratense</i>
Clover, white,	<i>Trifolium repens</i>
Cogongrass,	<i>Imperata cylindrica</i>
Dallisgrass,	<i>Paspalum dilatatum</i>
Dandelion,	<i>Taraxacum officinale</i>
Dock, curly,	<i>Rumex crispus</i>
Dogbane, hemp,	<i>Apocynum cannabinum</i>
Fescues,	<i>Festuca spp.</i>
Fescue, tall,	<i>Festuca arundinacea</i>
Guineagrass,	<i>Panicum maximum</i>
Horsenettle,	<i>Solanum carolinense</i>
Horseradish,	<i>Armoracia rusticana</i>

Ice plant,
Johnsongrass,
Kikuyugrass,
Knapweed,
Lantana,
Lespedeza,
Milkweed,
Muhly, wirestem,
Mullein, common,
Napiergrass,
Nightshade, silverleaf,
Nutsedge; purple, yellow,

Orchardgrass,
Pampasgrass,
Paragrass,
Phragmites*,
Poison hemlock,
Quackgrass,
Redvine*,
Reed, giant,
Ryegrass, perennial,
Smartweed, swamp,
Spurge, leafy*,
Starthistle, yellow,
Sweet potato, wild*,
Thistle, Canada,
Thistle, artichoke,
Timothy,
Torpedograss*,
Trumpet creeper*,
Vaseygrass,
Velvetgrass,
Wheatgrass, western,

Mesembryanthemum crystallinum
Sorghum halepense
Pennisetum clandestinum
Centaurea repens
Lantana camara
Lespedeza spp.
Asclepias spp.
Muhlenbergia frondosa
Verbascum thapsus
Pennisetum purpureum
Solanum elaeagnifolium
Cyperus rotundus,
Cyperus esculentus
Dactylis glomerata
Cortaderia spp.
Brachiaria mutica
Phragmites spp.
Conium maculatum
Agropyron repens
Brunnichia ovata
Arundo donax
Lolium perenne
Polygonum coccineum
Euphorbia esula
Centaurea solstitialis
Ipomoea pandurata
Cirsium arvense
Cynara cardunculus
Phleum pratense
Panicum repens
Campsis radicans
Paspalum urvillei
Holcus spp.
Agropyron smithii

*Partial Control

This product is not registered in California for use on water bermudagrass.

See "DIRECTIONS FOR USE" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for labeled uses and specific application instructions.

Alfalfa: Apply 1 quart of this product per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Make application after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.

Alligatorweed: Apply 4 quarts of this product per acre or apply a 1.5 percent solution with hand-held equipment to provide partial control. Apply when most of the plants are in bloom. Repeat applications will be required to maintain such control.

Anise (fennel)/Poison hemlock: Apply a 1 to 2 percent solution of this product as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Repeat applications may be needed in succeeding years to control plants arising from seeds.

Bentgrass: For suppression in grass seed production areas. For ground applications only, apply 1.5 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 20 gallons of water per acre. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should be actively growing and have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is recommended for best results. Failure to use tillage after treatment may result in unacceptable control.

Bermudagrass: For control, apply 5 quarts of this product per acre. For partial control, apply 3 quarts per acre. Treat when bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control. Allow 7 or more days after application before tillage.

Bermudagrass, water (knotgrass): Apply 1.5 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Apply when water bermudagrass is actively growing and 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field.

Fall applications only: Apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Fallow fields should be tilled prior to application. Apply prior to frost on water bermudagrass that is actively growing and 12 to 18 inches in length. Allow 7 or more days before tillage.

Bindweed, Field: For control, apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts east of the Mississippi River. Apply when the weeds are actively growing and are at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.

Also for control, apply 2 quarts of this product plus 0.5 pound a.i. of "Banvel" in 10 to 20 gallons of water per acre. At these rates, apply using ground application only.

The following tank mixtures with 2,4-D may be applied using aerial application equipment (except in California) in fallow and reduced tillage systems only.

For suppression on irrigated agricultural land, apply 1 to 2 quarts of this product plus 1 pound a.i. of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth.

For suppression, apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

In California only, apply 1 to 5 quarts of this product per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions.

For suppression on irrigated land where annual tillage is performed, apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to actively growing bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth. Allow 3 or more days after application before tillage.

Bluegrass, Kentucky/Bromegrass, smooth/Orchardgrass: Apply 2 quarts of this product in 10 to 40 gallons of water per acre when the grasses are actively growing and most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height. Allow 7 or more days after application before tillage.

Orchardgrass (sods going to no-till corn): Apply 1 to 1.5 quarts of this product per acre plus 0.5 to 1 percent non-ionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.

Blueweed, Texas: Apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when weed is actively growing and is at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.

Brackenfern: Apply 3 to 4 quarts of this product per acre as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 18 inches long.

Bursage, woollyleaf: For control, apply 2 quarts of this product plus 1 pint of "Banvel" per acre. For partial control, apply 1 quart of this product plus 1 pint of "Banvel" per acre. Add 0.5 to 1 percent nonionic surfactant by total spray volume and apply in 3 to 20 gallons of water per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.

Canarygrass, reed/Timothy/Wheatgrass, western: Apply 2 to 3 quarts of this product per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage.

Cogongrass: Apply 3 to 5 quarts of this product plus 0.5 to 1 percent nonionic surfactant in 10 to 40 gallons of water per acre. Apply when Cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

Dandelion/Dock, curly: Apply 3 to 5 quarts of this product per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.

Also for control, apply 16 fluid ounces of this product plus 0.5 pound a.i. 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre.

Dogbane, Hemp: Apply 4 quarts of this product per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.

For suppression, apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.

Fescue, Tall: Apply 3 quarts of this product in 10 to 40 gallons of water per acre to actively growing plants when most have reached boot-to-early seedhead stage of development.

Fall applications only--Apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to fescue in the fall when actively growing and plants have 6 to 12 inches of new growth. Allow 7 or more days after application before tillage. A sequential application of 1 pint per acre of this product plus nonionic surfactant will improve long-term control and control seedlings germinating after fall treatments or the following spring.

Guineagrass: Apply 3 quarts of this product per acre or use a 1 percent solution with hand-held equipment. Apply to actively growing guineagrass when most has reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. Allow 7 or more days after application before tillage.

Johnsongrass/Ryegrass, Perennial: Apply 1 to 3 quarts of this product per acre. In annual cropping systems apply 1 to 2 quarts of this product per acre. Apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. In noncrop, or areas where annual tillage (no-till) is not performed, apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tankmix with residual herbicides when using the 1 quart per acre rate.

For burndown of Johnsongrass, apply 1 pint per acre plus 0.5 to 1 percent nonionic surfactant in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.

Spot treatment (partial control or suppression)--Apply a 1 percent solution of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.

Kikuyugrass: Apply 2 to 3 quarts of this product per acre. Spray when most kikuyugrass is at least 8 inches in height (3 or 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.

Knapweed/Horseradish: Apply 4 quarts of this product per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.

Lantana: Apply this product as a 1 to 1.25 percent solution using hand-held equipment only. Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth. Allow 7 or more days after application before tillage.

Milkweed, Common: Apply 3 quarts of this product per acre. Apply when actively growing and most of the milkweed has reached the late bud to flower stage of growth. Following small grain harvest or mowing, allow milkweed to regrow to a mature stage prior to treatment. Allow 7 or more days after application before tillage.

Muhly, Wirestem: Apply 1 to 2 quarts of this product per acre. Use 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre or in pasture, sod, or noncrop areas. Spray when the wirestem muhly is 8 inches or more in height and actively growing. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage. This product will not provide residual control of wirestem muhly from seeds which germinate after application of this product. Do not tank mix with residual herbicides when using the 1 quart per acre rate.

Nightshade, Silverleaf: For control, apply 2 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Applications should be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth.

Nutsedge; Purple, Yellow: Apply 3 quarts of this product per acre as a broadcast spray, or apply a 1 to 2 percent solution from hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control of ungerminated tubers.

Sequential applications of 1 to 2 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre will provide control. Make applications when a majority of the plants are in the 3 to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3 to 5-leaf stage. Subsequent applications will be necessary for long-term control.

For suppression to partial control of existing plants, apply 1 pint to 2 quarts of this product per acre, plus 0.5 to 1 percent nonionic surfactant in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants. Wait 7 days after treatment before tillage or mowing.

Pampasgrass/Ice plant: Apply this product as a 1.5 to 2 percent solution using hand-held equipment. Apply to plants that are actively growing. Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

Phragmites: For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 5 quarts per acre as a broadcast spray or apply a 2 percent solution from hand-held equipment. In other areas of the U.S., apply 3 quarts per acre as a broadcast spray or apply a 1 percent solution from hand-held equipment for partial control. For best results, treat during late summer or fall months or when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.

Quackgrass--In Annual Cropping Systems, or in Pastures and Sods Followed by Deep Tillage: Apply 1 to 2 quarts of this product per acre. For the 1 quart rate, apply 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. For the 2 quart rate, apply in 10 to 40 gallons of water per acre. Do not tank mix with residual herbicides when using the 1 quart rate. Spray when quackgrass is 6 to 8 inches in height and actively growing. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, for best results use a moldboard plow.

Quackgrass--Pasture or Sod or Other Noncrop Areas Where Deep Tillage is Not Planned Following Application: Apply 2 to 3 quarts in 10 to 40 gallons of water per acre. Spray when the quackgrass is greater than 8 inches tall and actively growing. Do not till between harvest and fall application or in fall or spring prior to spring application. Allow 3 or more days after application before tillage.

Redvine: For suppression, apply 24 fluid ounces of this product per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply recommended rates in 5 to 10 gallons of water per acre plus 0.5 to 1 percent nonionic surfactant by total volume. Apply in late September or early October to actively growing plants, which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Reed, Giant: For control of giant reed, apply a 2 percent solution of this product when plants are actively growing. Best results are obtained when applications are made in late summer to fall.

Smartweed, Swamp: Apply 3 to 5 quarts of this product per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.

Also for control, apply 16 fluid ounces of this product plus 0.5 pound active ingredient of 2,4-D plus 0.5 to 1 percent nonionic surfactant by total volume in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.

Spurge, Leafy: For suppression, apply 16 fluid ounces of this product plus 0.5 pound active ingredient 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall. Allow 7 or more days after application before tillage.

Starthistle, Yellow: Best results are obtained when applications are made during periods of active growth, including the rosette, bolting and early flower stages. For spray-to-wet applications, apply this product as a 2 percent solution. For broadcast applications, apply 2 quarts per acre in 10 to 40 gallons per acre of water carrier.

Sweet Potato, Wild/Thistle, Artichoke: Apply this product as a 2 percent solution using hand-held equipment. Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications may be required. Allow the plant to reach the recommended stage of growth before retreatment. Allow 7 or more days before tillage.

Thistle, Canada: Apply 2 to 3 quarts of this product per acre. Apply to actively growing thistles when most are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

For suppression of Canada thistle, apply 1 quart per acre of this product, or 1 pint of this product plus 0.5 pound a.i. 2,4-D per acre, plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating.

Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.

Torpedograss: Apply 4 to 5 quarts of this product per acre to provide partial control of torpedograss. Apply to actively growing torpedograss when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost. Allow 7 or more days after application before tillage.

Trumpet creeper: For control, apply 2 quarts of this product per acre in 5 to 10 gallons of water per acre. Apply to actively growing plants in late September or October, which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Other perennials listed on this label: Apply 3 to 5 quarts of this product per acre. Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.

WOODY BRUSH AND TREES

When applied as recommended under the conditions described, this product CONTROLS or PARTIALLY CONTROLS the following woody brush, plants and trees:

Alder,	<i>Alnus spp.</i>
Ash*,	<i>Fraxinus spp.</i>
Aspen, quaking,	<i>Populus tremuloides</i>
Bearmat (Bearclover),	<i>Chamaebatia foliolosa</i>
Beech,	<i>Fagus grandifolia</i>
Birch,	<i>Betula spp.</i>
Blackberry,	<i>Rubus spp.</i>
Blackgum,	<i>Nyssa spp.</i>
Bracken,	<i>Peridium spp.</i>
Broom:	
French,	<i>Cytisus monspessulanus</i>
Scotch,	<i>Cytisus scoparius</i>
Buckwheat, California*,	<i>Eriogonum fasciculatum</i>
Cascara*,	<i>Rhamnus purshiana</i>
Catsclaw*,	<i>Acacia greggi</i>
Ceanothus*,	<i>Ceanothus spp.</i>
Chamise,	<i>Adenostoma fasciculatum</i>

Cherry:	
Bitter,	<i>Prunus emarginata</i>
Black,	<i>Prunus serotina</i>
Pin,	<i>Prunus pensylvanica</i>
Coyote brush,	<i>Baccharis consanguinea</i>
Creeper, Virginia*,	<i>Parthenocissus quinquefolia</i>
Dewberry,	<i>Rubus trivialis</i>
Dogwood*,	<i>Cornus spp.</i>
Elderberry,	<i>Sambucus spp.</i>
Elm*,	<i>Ulmus spp.</i>
Eucalyptus,	<i>Eucalyptus spp.</i>
Gorse,	<i>Ulex europaeus</i>
Hasardia*,	<i>Haplopappus squamosus</i>
Hawthorn,	<i>Crataegus spp.</i>
Hazel,	<i>Corylus spp.</i>
Hickory*,	<i>Carya spp.</i>
Holly, Florida/Brazilian Peppertree*,	<i>Schinus terebinthifolius</i>
Honeysuckle,	<i>Lonicera spp.</i>
Hornbeam, American*,	<i>Carpinus caroliniana</i>
Kudzu,	<i>Pueraria lobata</i>
Locust, black*,	<i>Robinia pseudoacacia</i>
Madrone,	<i>Arbutus menziesii</i>
Manzanita,	<i>Arctostaphylos spp.</i>
Maple:	
Red**,	<i>Acer rubrum</i>
Sugar,	<i>Acer saccharum</i>
Vine*,	<i>Acer circinatum</i>
Monkey Flower*,	<i>Mimulus guttatus</i>
Oak:	
Black*,	<i>Quercus velutina</i>
Northern Pin,	<i>Quercus palustris</i>
Post,	<i>Quercus stellata</i>
Red,	<i>Quercus rubra</i>
Southern Red,	<i>Quercus falcata</i>
White*,	<i>Quercus alba</i>
Persimmon*,	<i>Diospyros spp.</i>
Pine,	<i>Pinus spp.</i>
Poison Ivy,	<i>Rhus radicans</i>
Poison Oak,	<i>Rhus toxicodendron</i>
Poplar, yellow*,	<i>Liriodendron tulipifera</i>
Raspberry,	<i>Rubus spp.</i>
Redbud, eastern,	<i>Cercis canadensis</i>
Rose, multiflora,	<i>Rosa multiflora</i>
Russian-olive,	<i>Elaeagnus angustifolia</i>
Sage; black, white,	<i>Salvia spp.</i>
Sagebrush, California,	<i>Artemisia californica</i>
Salmonberry,	<i>Rubus spectabilis</i>
Salt cedar,	<i>Tamarix spp.</i>
Sassafras,	<i>Sassafras aioidum</i>
Sourwood,	<i>Oxydendrum arboreum</i>
Sumac:	
Poison*,	<i>Rhus vernix</i>
Smooth*,	<i>Rhus glabra</i>
Winged*,	<i>Rhus copallina</i>
Sweetgum,	<i>Liquidambar styraciflua</i>
Swordfern*,	<i>Polystichum munitum</i>
Tallowtree, Chinese,	<i>Sapium sebiferum</i>
Tan Oak,	<i>Lithocarpus densiflorus</i>
Thimbleberry,	<i>Rubus parviflorus</i>
Tobacco, tree*,	<i>Nicotiana glauca</i>
Trumpetcreeper,	<i>Campsis radicans</i>
Waxmyrtle, southern*,	<i>Myrica cerifera</i>
Willow,	<i>Salix spp.</i>

* Partial control

** See below for control or partial control instructions.

NOTE: If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stages of growth.

Apply this product when plants are actively growing and, unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

See "DIRECTIONS FOR USE" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for labeled uses and specific application instructions.

Apply this product as follows to control or partially control the following woody brush and trees.

Alder/Dewberry/Honeysuckle/Post Oak/Raspberry: For control, apply 3 to 4 quarts per acre of this product as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment.

Aspen, Quaking/Cherry:Bitter, Black, Pin/Hawthorn /Oak, Southern Red/Sweetgum/Trumpetcreeper: For control, apply 2 to 3 quarts of this product per acre as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment.

Birch/Elderberry/Hazel/Salmonberry/Thimbleberry: For control, apply 2 quarts per acre of this product as a broadcast spray or as a 1 percent solution with hand-held equipment.

Blackberry: For control, apply 3 to 4 quarts per acre of this product as a broadcast spray, or 1 to 1.5 percent solution with hand-held equipment. Make application after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. After berries have set or dropped in late fall, blackberry can be controlled by applying a 3/4 percent solution of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume with hand-held equipment. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of this product in 10 to 40 gallons of water per acre.

Broom: French, Scotch: For control, apply a 1.5 to 2 percent solution with hand-held equipment.

Buckwheat, California/Hasardia/Monkey Flower/Tobacco, Tree: For partial control of these species, apply a 1 to 2 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Catsclaw: For partial control, apply as a 1 to 1.5 percent solution with hand-held equipment.

Coyote Brush: For control, apply a 1.5 to 2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Eucalyptus: For control of eucalyptus resprouts, apply a 2 percent solution of this product with hand-held equipment when resprouts are 6 to 12 feet tall. Ensure complete coverage. Apply when plants are growing actively. Avoid application to drought-stressed plants.

Kudzu: For control, apply 4 quarts of this product per acre as a broadcast spray or as a 2 percent solution with hand-held equipment. Repeat applications will be required to maintain control.

Madrone resprouts: For suppression or partial control, apply a 2 percent solution of this product to resprouts less than 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.

Maple, Red: For control, apply as a 1 to 1.5 percent solution with hand-held equipment when at least 50% of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of this product per acre as a broadcast spray.

Maple, Sugar/Oak, Northern Pin/Oak, Red: For control, apply as a 1 to 1.5 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Poison Ivy/Poison Oak: For control, apply 4 to 5 quarts of this product per acre as a broadcast spray or as a 2 percent solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.

Rose, Multiflora: For control, apply 2 quarts of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Treatments should be made prior to leaf deterioration by leaf-feeding insects.

Sage, Black/Sagebrush, California/Chamise/Tallowtree, Chinese: For control of these species, apply a 1 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Tan Oak Resprouts: For suppression or partial control, apply a 2 percent solution of this product to resprouts less than 3 to 6 feet tall. Best results are obtained with fall applications.

Willow: For control, apply 3 quarts of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment.

Other Woody Brush and Trees listed on this label: For partial control, apply 2 to 5 quarts of this product per acre as a broadcast spray or as a 1 to 2 percent solution with hand-held equipment.

NONCROP USES

See "GENERAL INFORMATION" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for essential product performance information and the following "NONCROP" sections for specific recommended uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE TURFGRASSES, TREES, SHRUBS OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

This product does not provide residual weed control. For subsequent weed control, follow a label-approved herbicide program.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

INDUSTRIAL, RECREATIONAL AND PUBLIC AREAS

When applied as directed for "NONCROP USES", under conditions described, this product controls annual and perennial weeds listed on this label growing in areas such as airports, ditch banks, dry ditches, dry canals, fencerows, golf courses, highways, industrial plant sites, lumber yards, parking areas, parks, petroleum tank farms and pumping installations, pipelines, power and telephone rights-of-way, railroads, roadsides, schools, storage areas, utility substations, other public areas and similar industrial or noncrop areas.

For specific rates of application and instructions for control of various annual and perennial weeds and woody brush and trees, see the "WEEDS CONTROLLED" section of this label.

This product may be applied with recirculating sprayers, shielded applicators, or wiper applicators in any noncrop site specified on this label. See the Selective Equipment part of "APPLICATION EQUIPMENT and TECHNIQUES" section of this label for information on proper use and calibration of this equipment.

TANK MIXTURES FOR INDUSTRIAL SITES AND FORESTRY SITE PREPARATIONS

GLYPHOSATE plus DuPont OUST®

Use on industrial sites including airports, industrial plants, lumberyards, petroleum tank farms, pumping stations, railroads, roadsides, storage areas or other similar sites where bare ground is desired.

This tank mixture may also be used as a site preparation treatment for sites to be planted to jack pine, loblolly pine, red pine, slash pine and Virginia pine. When applied as directed for "NONCROP USES" under the conditions described, this product plus OUST® provides control of annual weeds listed in the "WEEDS CONTROLLED" section of the label for this product and OUST®, and partial control of the perennial weeds listed below.

Apply 1 to 2 quarts of this product with 2 to 4 ounces of OUST® in 10 to 40 gallons of spray solution per acre as a broadcast spray to actively growing weeds.

This mixture may be applied by aerial equipment in site prep operations. When applied by air, use the recommended rates in 5 to 15 gallons of spray solution per acre.

This product plus OUST® tank mixtures may not be applied by air in California.

For control of annual weeds, use the lower rates of these products.

For control of the listed perennial weeds, use the higher rates of both products. For partial control, use the lower rates.

Bahiagrass, *Paspalum notatum*

Bermudagrass*, *Cynodon dactylon*

Broomsedge, *Andropogon virginicus*

Dock, curly, *Rumex crispus*

Dogfennel, *Eupatorium capilliflorum*

Fescue, tall, *Festuca arundinacea*

Johnsongrass**, *Sorghum halepense*

Poorjoe**, *Diodia teres*

Quackgrass, *Agropyron repens*

Trumpetcreeper*, *Campsis radicans*

Vaseygrass, *Paspalum urvillei*

Vervain, blue, *Verbena hastata*

*Suppression at higher rates only.

**Control at lower rates.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

TANK MIXTURES FOR NONCROP SITES

When applied as a tank mixture, this product provides control of the emerged annual weeds and partial control of the emerged perennial weeds listed in this label. When applied as a tank mixture, the following residual herbicides will provide preemergence control of the weeds listed in the individual product labels.

GLYPHOSATE may be tank mixed with products such as: DuPont KROVAR® I DF, ESCORT®, TELAR®, OUST®, VELPAR®, “Diuron 4L”, “Karmex DF”, “Surflan”, “Stomp”.

When tank mixing with residual herbicides, add an agriculturally approved nonionic surfactant at 0.5 to 1 percent by volume of spray solution. See the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label before preparing these tank mixtures.

Read and carefully observe the label claims, cautionary statements, recommended use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

CONTROL OF EMERGED WEEDS

Note: For backpack sprayer and handgun applications, see the "HAND-HELD AND HIGH VOLUME EQUIPMENT" section for recommended rates.

Annual Weeds: Apply 1 quart per acre of this product in these tank mixtures when weeds are less than 6 inches tall and 1.5 quarts per acre when weeds are more than 6 inches tall.

Perennial Weeds: For partial control of perennial weeds using these tank mixtures, apply 2 to 5 quarts per acre of this product. Follow the recommendations in the "WEEDS CONTROLLED" section of this label for stage of growth and rate of application for specific perennial weeds.

PREEMERGENCE WEED CONTROL

For preemergence weed control, refer to the individual product labels for specific noncrop sites, rates, carrier volumes and precautionary statements.

Mix only the quantity of spray solution which can be used during the same day. Do not allow these tank mixtures to stand overnight as this may result in reduced weed control.

FARMSTEAD WEED CONTROL

When applied as directed for "NONCROP USES", under conditions described, this product controls undesirable vegetation listed on this label around farmstead building foundations, along and in fences, shelterbelts and for general nonselective farmstead weed control.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

FARM DITCHES

This product will suppress perennial grasses along farm ditches. Apply this product at a rate of 6 to 8 fluid ounces per acre. Use 8 fluid ounces per acre when treating tall (coarse) fescue, fine fescue, orchardgrass or quackgrass covers. For best suppression of these species, add ammonium sulfate at a rate of 1.7 pounds per 10 gallons of spray solution. Use 6 fluid ounces per acre without ammonium sulfate when treating Kentucky bluegrass.

Apply treatments in 10 to 20 gallons of spray solution per acre to actively growing perennial grass covers. For best spray distribution and coverage, use flat fan nozzles.

Add a nonionic surfactant at a rate of 0.5 percent of the spray solution.

Where broadleaf weed control or suppression is desired, tank mix this product with an appropriate, labeled broadleaf weed herbicide.

CONSERVATION RESERVE PROGRAM (CRP ACRES)

This product can be used to control undesirable vegetation when rotating out of CRP acres or to suppress competitive growth and seed production of undesirable vegetation in CRP acres.

For specific rates of application for various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

CRP applications may be made with wiper applicators or conventional spray equipment.

For selective applications with broadcast spray equipment, apply 12 to 16 ounces per acre of this product in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy. Some stunting of CRP perennial grasses will occur if applications are made when plants are not dormant.

DORMANT RANGELAND*

This product will control or suppress many weeds, including downy brome, cheat grass, cereal rye, medusahead rye and jointed goatgrass in dormant rangeland.

Apply 8 to 16 ounces per acre of this product in the early spring when the weeds have greened up, but desirable grasses, such as crested and tall wheatgrass are still truly dormant.

Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off.

Do not use additional surfactant or ammonium sulfate when spraying dormant rangeland grasses with GLYPHOSATE.

*In California, restrict applications to less than 10% of dormant range and acreage. Make spot treatments only.

HABITAT MANAGEMENT

This product is recommended for the restoration and/or maintenance of native habitats and in wildlife management areas. Apply as recommended in the "NONCROP USES" section of this label.

Habitat Restoration and Maintenance: When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broadspectrum vegetation control requirements in habitat management areas. Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement. For spot treatments, care should be exercised to keep spray off of desirable plants.

Wildlife Food Plots: This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after applying this product before tilling.

CHRISTMAS TREES

THIS PRODUCT IS NOT RECOMMENDED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY.

Note: Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

When applied as instructed for the conditions described for "NONCROP USES", this product controls undesirable vegetation listed on this label prior to planting as a postdirected spray around Christmas trees.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

Site Preparation: Following preplant applications of this product, Christmas tree species may be planted. Precautions should be taken to protect nontarget plants during site preparation applications.

Postdirected Spray: Use as a postdirected spray around Christmas trees such as those listed below. Care must be exercised to avoid contact of spray, drift or mist with foliage of or green.

Fir, Abies spp.,	<i>Pseudotsuga spp.</i>
Pine,	<i>Pinus spp.</i>
Spruce,	<i>Picea spp.</i>

SILVICULTURAL SITES AND RIGHTS-OF-WAY

NOTE: NOT RECOMMENDED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN SILVICULTURAL NURSERIES

When applied as directed for "NONCROP USES" under conditions described, this product controls undesirable vegetation listed on this label. This product also suppresses or controls undesirable vegetation listed on this label when applied at recommended rates for release of established coniferous species listed on this label.

For specific rates of application and instructions for control of various brush, annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label. For specific rates of application for release of listed coniferous species, see the "CONIFER RELEASE" part of this section of the label.

Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

Aerial Application: This product may be applied using aerial spray equipment for silvicultural site preparation, conifer release and rights-of-way treatments. See the "APPLICATION EQUIPMENT and TECHNIQUES" part of the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label for information on how to apply this product by air.

DO NOT APPLY THIS PRODUCT BY AIR TO RIGHTS-OF-WAY SITES IN THE STATE OF CALIFORNIA.

SITE PREPARATION

Following preplant applications of this product, any silvicultural species may be planted.

POSTDIRECTED SPRAY

In established silvicultural sites, use as a spray on the foliage of undesirable vegetation. Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of desirable species.

CONIFER RELEASE

For release, apply only where conifers have been established for more than one year. Vegetation should not be disturbed prior to treatment or until visual symptoms appear after treatment. Symptoms of treatment are slow to appear, especially in woody species treated in late fall. Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied or when applications are made during periods of active conifer growth. **Do not use additional surfactant with conifer release applications.**

Applications must be made after formation of final conifer resting buds in the fall or prior to initial bud swelling in spring. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Use the following rates for conifer release to control or partially control the weeds listed in the "WEEDS CONTROLLED" section of this label.

For release of the following conifer species:

Douglas fir,	<i>Pseudotsuga menziesii</i>
Fir,	<i>Abies spp.</i>
Hemlock,	<i>Tsuga spp.</i>
Pines*,	<i>Pinus spp.</i>
Spruce,	<i>Picea spp.</i>

* Includes all species except eastern white pine, loblolly pine or slash pine.

Apply 1.5 to 2 quarts of this product per acre except in Washington and Oregon, west of the crest of the Cascade Mountains. For spring treatments west of the crest of the Cascade Mountains, apply 1 quart of this product per acre before conifer bud swell for control of annual weeds. For fall treatments in Washington and Oregon, west of the crest of the Cascade Mountains, apply 1 to 1.5 quarts of this product per acre before any major leaf drop of deciduous species.

For release of western hemlock, apply 1 quart of this product per acre.

For release of the following conifer species:

Loblolly pine,	<i>Pinus taeda</i>
Eastern white pine,	<i>Pinus strobus</i>
Slash pine,	<i>Pinus elliottii</i>

Late Season Application - Apply 1.5 to 2 quarts of this product in a minimum of 5 gallons of spray solution per acre during early autumn. Applications made prior to September 1 or when conditions are conducive to rapid growth of conifers will create the potential for increased injury in the form of tip and/or needle burn. Injury may decrease with later applications. Some autumn colors are acceptable at time of application. Apply prior to frost or leaf drop of undesirable plants. Applications made according to label directions will release loblolly pine, eastern white pine and slash pine by reducing competition from the following species:

Ash,	<i>Fraxinus spp.</i>
Cherry:	
Black,	<i>Prunus serotina</i>
Pin,	<i>Prunus pensylvanica</i>
Elm,	<i>Ulmus spp.</i>
Hawthorn,	<i>Crataegus spp.</i>
Locust, black,	<i>Robina pseudoacacia</i>
Maple, red,	<i>Acer rubra</i>
Oak:	
Black,	<i>Quercus velutina</i>
Post,	<i>Quercus stellata</i>
Southern Red,	<i>Quercus falcata</i>
White,	<i>Quercus alba</i>
Persimmon,	<i>Diospyros spp.</i>
Poplar, yellow,	<i>Liriodendron tulipifera</i>
Sassafras,	<i>Sassafras albidum</i>
Sourwood,	<i>Oxydendrum arboreum</i>
Sumac:	
Poison,	<i>Rhus vernix</i>
Smooth,	<i>Rhus glabra</i>
Winged,	<i>Rhus copallina</i>
Sweetgum,	<i>Liquidambar styraciflua</i>

Apply only to those sites where woody brush and trees listed in this label constitute the majority of the undesirable species.

GLYPHOSATE plus DuPont OUST® Tank Mixtures for Conifer Release from Herbaceous Weeds

To release loblolly pines from herbaceous weeds, tank mixtures of this product with OUST® will provide control of annual weeds listed in the "WEEDS CONTROLLED" section of this and the OUST® label, and partial control of the perennial weeds listed below.

Apply 16 to 24 fluid ounces of this product with 2 to 4 ounces of OUST® in 10 to 30 gallons of spray solution per acre. Make application to actively growing weeds as a broadcast spray over the top of the young loblolly pines.

This product plus OUST® tank mixtures may not be applied by air in California.

This tank mixture may be applied using aerial equipment. When applying by air, use the recommended rate in 5 to 15 gallons of spray solution per acre.

For control of annual weeds below 12 inches in height (or runner length on annual vines), use the lower rates of both products. Use the higher rates of both products when annual weeds are in more advanced stages of growth and approaching flower or seed formation.

Use the higher rates of both products for partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass,	<i>Paspalum notatum</i>
Broomsedge,	<i>Andropogon virginicus</i>
Dock, curly,	<i>Rumex crispus</i>
Dogfennel,	<i>Eupatorium capilliflorum</i>
Fescue, tall,	<i>Festuca arundinacea</i>
Johnsongrass*,	<i>Sorghum halepense</i>
Poorjoe*,	<i>Diodia teres</i>
Trumpet creeper**,	<i>Campsis radicans</i>
Vaseygrass,	<i>Paspalum urvillei</i>
Vervain, blue,	<i>Verbena hastata</i>

* Control at the higher rates.

** Suppression at the higher rates only.

Pine damage may occur or can be accentuated if treatment takes place when young trees are under stress from drought, flood water, insects or disease.

Read and observe the cautionary statements and all other information appearing on the labels of all herbicides used.

NOTE TO USER: This product must not be used in areas where adverse impact on federally designated endangered/threatened plant or aquatic species is likely.

Prior to making applications, the user of this product must determine that no such species are located in or immediately adjacent to the area to be treated.

CUT STUMP TREATMENTS

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly-cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

When used according to directions for cut stump application, this product will CONTROL, PARTIALLY CONTROL or SUPPRESS many types of woody brush and tree species, some of which are listed below:

Alder,	<i>Alnus spp.</i>
Eucalyptus,	<i>Eucalyptus spp.</i>
Madrone,	<i>Arbutus menziesii</i>
Oak,	<i>Quercus spp.</i>
Reed, giant,	<i>Arundo donax</i>
Saltcedar,	<i>Tamarisk spp.</i>
Sweetgum,	<i>Liquidambar styraciflua</i>
Tan Oak,	<i>Lithocarpus densiflorus</i>
Willow,	<i>Salix spp.</i>

INJECTION AND FRILL APPLICATIONS

Woody vegetation may be controlled by injection or frill application of this product. Apply this product using suitable equipment which must penetrate into the living tissue. Apply the equivalent of 1 ml of this product per each 2 to 3 inches of trunk diameter (DBH). This is best achieved by applying a 50 to 100 percent concentration of this material either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In species such as this, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, application should be made during periods of active growth and after full leaf expansion.

This treatment WILL CONTROL the following woody species:

Oak,	<i>Quercus spp.</i>
Poplar,	<i>Populus spp.</i>
Sweetgum,	<i>Liquidambar styraciflua</i>
Sycamore,	<i>Platanus occidentalis</i>

This treatment WILL SUPPRESS the following woody species:

Black gum,	<i>Nyssa sylvatica</i>
Dogwood,	<i>Cornus spp.</i>
Hickory,	<i>Carya spp.</i>
Maple, red,	<i>Acer rubrum</i>

TURFGRASSES AND GRASSES FOR SEED PRODUCTION

PREPLANT AND RENOVATION

When applied as directed for "NONCROP USES", under conditions described, this product controls most existing vegetation prior to the planting or renovation of either turfgrasses or grass seed production areas.

For specific rates of application and instructions for control of various annual and perennial weeds, and woody brush and trees, see the "WEEDS CONTROLLED" section of this label.

For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as bermudagrass, summer or fall applications provide best control.

DO NOT DISTURB SOIL OR UNDERGROUND PLANT PARTS BEFORE TREATMENT. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.

TURFGRASSES

Where existing vegetation is growing in a field or unmowed situation, apply this product to actively growing weeds at the stages of growth listed in the "WEEDS CONTROLLED" section of this label.

Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray.

Desirable turfgrasses may be planted following the above procedures.

GRASSES FOR SEED PRODUCTION

Apply this product to actively growing weeds at the stages of growth recommended in the "WEEDS CONTROLLED" section of this label prior to planting or renovation of turf or forage grass areas grown for seed production.

DO NOT feed or graze treated areas within 8 weeks after application.

ANNUAL WEED CONTROL IN DORMANT BERMUDAGRASS AND BAHIAGRASS TURF

When applied as directed for "NONCROP USES" under the conditions described, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass turf. Refer to the rate table for GLYPHOSATE alone under the "RELEASE OF BERMUDAGRASS and BAHIAGRASS" section of this label for recommended rates and volumes on the species to be suppressed or controlled. Treat only when turf is dormant and prior to spring greenup. Spot treatments or broadcast applications of this product in excess of 16 fluid ounces per acre may result in injury or delayed greenup in highly maintained turfgrass areas; i.e., golf courses, lawns, etc. DO NOT APPLY TANK MIXTURES of this product plus OUST® in highly maintained turfgrass areas.

RELEASE OF BERMUDAGRASS OR BAHIAGRASS

NOTE: Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. Use tank mixtures of this product plus OUST® only on railroads, highways, utility plant sites, or other right-of-way areas.

When applied as directed for "NONCROP USES" under the conditions described, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. This product may be tankmixed with OUST® as recommended for residual control. Make applications to dormant bermudagrass or bahiagrass. Tank mixtures of this product plus OUST® may delay greenup. To avoid delays in greenup and minimize injury, do not add more than 1 ounce per acre of OUST® on bermudagrass or more than 0.5 ounce per acre on bahiagrass, or treat when these grasses are in a semi-dormant condition.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4 to 6-leaf stage.

WEEDS CONTROLLED

Rate recommendations for control or suppression of winter annuals and tall fescue are listed below:

Apply the recommended rates of this product alone or as a tank mixture in 10 to 25 gallons of water, plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre.

For the best recommendation for the mixture of weeds within your geographic area, contact your Dupont sales representative.

WEEDS CONTROLLED OR SUPPRESSED WITH GLYPHOSATE ALONE*

NOTE: C = Control
S = Suppression

WEED SPECIES	GLYPHOSATE FLUID OZ/ACRE					
	8	12	16	24	32	64
Barley, little, <i>Hordeum pusillum</i>	S	C	C	C	C	C
Bedstraw, catchweed <i>Galium aparine</i>	S	C	C	C	C	C
Bluegrass, annual <i>Poa annua</i>	S	C	C	C	C	C
Chervil <i>Chaerophyllum tainturieri</i>	S	C	C	C	C	C
Chickweed, common <i>Stellaria media</i>	S	C	C	C	C	C
Clover, crimson <i>Trifolium incarnatum</i>	•	S	S	C	C	C
Clover, large hop <i>Trifolium campestre</i>	•	S	S	C	C	C
Fescue, tall <i>Festuca arundinaceae</i>	•	•	•	•	S	S
Geranium, Carolina <i>Geranium carolinianum</i>	•	•	S	S	C	C
Henbit <i>Lamium amplexicaule</i>	•	S	C	C	C	C
Ryegrass Italian <i>Lolium multiflorum</i>	•	•	S	C	C	C
Speedwell, corn <i>Veronica arvensis</i>	S	C	C	C	C	C
Vetch, common <i>Vicia sativa</i>	•	•	S	C	C	C

* These rates apply only to sites where an established competitive turf is present.

**WEEDS CONTROLLED OR SUPPRESSED
WITH GLYPHOSATE PLUS DUPONT OUST®**

NOTE: C = Control
S = Suppression

GLYPHOSATE + OUST®								
WEED SPECIES	GLYPHOSATE (FL OZ/A)	8	12	12	16	16	12	16
	OUST® (OZ/A)	1/4	1/4	1/2	1/4	1/2	1	1
Barley, little								
<i>Hordeum pusillum</i>		C	C	C	C	C	C	C
Bedstraw, catchweed								
<i>Galium aparine</i>		C	C	C	C	C	C	C
Bluegrass, annual								
<i>Poa annua</i>		S	C	C	C	C	C	C
Chervil								
<i>Chaerophyllum tainturieri</i>		C	C	C	C	C	C	C
Chickweed, common								
<i>Stellaria media</i>		S	C	C	C	C	C	C
Clover, crimson								
<i>Trifolium incarnatum</i>		S	S	S	S	C	C	C
Clover, large hop								
<i>Trifolium campestre</i>		•	•	S	S	S	C	C
Fescue, tall								
<i>Festuca arundinacea</i>		•	•	•	•	•	S	S
Geranium, Carolina								
<i>Geranium carolinianum</i>		•	S	S	C	C	C	C
Henbit								
<i>Lamium amplexicaule</i>		•	S	C	C	C	C	C
Ryegrass, Italian								
<i>Lolium multiflorum</i>		•	S	S	C	C	C	C
Speedwell, corn								
<i>Veronica arvensis</i>		S	C	C	C	C	C	C
Vetch, common								
<i>Vicia sativa</i>		C	C	C	C	C	C	C

* These rates or mixtures of rates apply only to sites where an established competitive turf is present.

**RELEASE OF ACTIVELY GROWING
BERMUDAGRASS**

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual species listed in the "WEEDS CONTROLLED" section of this and the OUST® label, and suppression or partial control of certain perennial weeds.

For control or suppression of those annual species listed on this label, use 1 to 3 pints of this product as a broadcast spray in 10 to 25 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation.

Use the higher rate of this product for partial control of the following perennial species. Use the lower rates for suppression of growth. For best results, see the "WEEDS CONTROLLED" section of this label for proper stage of growth.

Bahiagrass,	<i>Paspalum notatum</i>
Bluestem, silver,	<i>Andropogon saccharoides</i>
Fescue, tall,	<i>Festuca arundinacea</i>
Johnsongrass**,	<i>Sorghum halepense</i>
Trumpetcreeper**,	<i>Campsis radicans</i>
Vaseygrass,	<i>Paspalum urvillei</i>

**Suppression at higher rates only.

This product may be tankmixed with OUST®. If tankmixed, use no more than 1 to 2 pints per acre of this product with 1 to 2 ounces of OUST® per acre.

Use the lower rates of both mixtures to control annual weeds below 6 inches in height (or runner length in annual vines) that are listed in the "WEEDS CONTROLLED" section of this booklet and the OUST® label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages.

Use the higher rates of this product to provide partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass,	<i>Paspalum notatum</i>
Bluestem, silver,	<i>Andropogon saccharoides</i>
Broomsedge,	<i>Andropogon virginicus</i>
Dock, curly,	<i>Rumex crispus</i>
Dogfennel,	<i>Eupatorium capilliflorum</i>
Fescue, tall,	<i>Festuca arundinacea</i>
Johnsongrass,	<i>Sorghum halepense</i>
Poorjoe,	<i>Diodia teres</i>
Trumpetcreeper,	<i>Campsis radicans</i>
Vaseygrass,	<i>Paspalum urvillei</i>
Vervin, blue,	<i>Verbena hastata</i>

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may result.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

COOL SEASON TURF GROWTH REGULATION

When applied as directed, this product will suppress growth and seedhead development of listed turf species in industrial sites.

This product is recommended for management of coarse turf on roadside rights-of-way or other industrial areas. Do not use on high-quality turf or other areas where some turf color changes cannot be tolerated. Slight turf discoloration may occur but turf will regreen and regrow under moist conditions as effects of this product wear off.

Apply 4 to 6 fluid ounces of this product per acre alone or in a recommended tank mixture. Spray volumes of 10 to 40 gallons per acre are recommended.

When using this product, mix 2 quarts of a nonionic surfactant per 100 gallons of spray solution.

This product can be used for growth and seedhead suppression of:

**Tall Fescue
Smooth Brome**

For best results, apply this product in a recommended tank mixture to actively growing turfgrasses after greenup in the spring of the year. For suppression of seedheads, applications must be made before boot-to-seedhead stage of development. Applications made from seedhead emergence until maturity may result in turf discoloration or injury.

After mowing or removal of seedheads, this product in a recommended tank mixture may also be used to suppress the growth of certain turfgrasses. Allow turf to recover from stress caused by heat, drought or mowing before making applications. Applications made to turf under stress may increase the potential for discoloration or injury.

ANNUAL GRASSES

For growth suppression of some annual grasses such as annual ryegrass, wild barley and wild oats, apply 3 to 4 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

TANK MIXTURES

For the following tank mixtures, consult each product label for weeds controlled and the correct stage of application. Do not treat turf under stress.

Tank mixtures plus 2,4-D Amine

For additional weed control benefits, up to 1 pound a.i. per acre of 2,4-D amine may be added to the following tank mixtures. Consult the label for 2,4-D amine for weeds controlled.

TALL FESCUE

GLYPHOSATE plus DuPont TELAR®

For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.5 ounce of TELAR® per acre.

This tank mixture can also be applied after mowing or removal of tall fescue seedheads for turf growth suppression. Make only one of the above applications per growing season.

GLYPHOSATE plus DuPont OUST®

For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.25 ounce of OUST® per acre.

GLYPHOSATE plus DuPont ESCORT®

This tank mixture can be applied after mowing or removal of tall fescue seedheads for turf growth suppression and control or partial control of some annual weeds. Use up to 1/3 ounce of ESCORT® per acre.

NOTE: This product is not registered for use with ESCORT® in California.

SMOOTH BROME

GLYPHOSATE plus OUST®

For suppression of smooth brome growth and seedheads and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.25 ounce of OUST® per acre.

BAHIAGRASS SEEDHEAD AND VEGETATIVE SUPPRESSION

When applied as directed in the indicated noncrop areas (roadsides, airports, golf course roughs, and plant sites), this product will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with single applications and approximately 120 days with sequential applications.

Apply this product 1 to 2 weeks after full greenup of bahiagrass or after the bahiagrass has been mowed to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 6 fluid ounces per acre of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 25 gallons of water per acre.

Sequential applications of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume may be made at approximately 45 day intervals to extend the period of seedhead and vegetative growth suppression. For continued seedhead suppression, sequential applications must be made prior to seedhead emergence. Apply no more than 2 sequential applications per year. As a first sequential application, apply 4 fluid ounces of this product per acre plus nonionic surfactant. A second sequential application of 2 to 4 fluid ounces per acre plus nonionic surfactant may be made approximately 45 days after the last application.

A tank mixture of this product plus OUST® may be applied only on roadsides for seedhead inhibition and vegetative suppression. Apply 6 fluid ounces per acre of this product plus 0.25 ounce per acre of OUST®, plus 0.5 to 1 percent nonionic surfactant by total spray volume 1 to 2 weeks following an initial spring mowing. When using this product plus OUST® for suppression of bahiagrass, make only 1 application per year.

CROPPING SYSTEMS

When applied as directed for "CROPPING SYSTEMS", under the conditions described, this product controls annual and perennial weeds listed on this label, prior to the emergence of direct seeded crops or prior to transplanting of crops listed on this label.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for essential product performance information.

See the following "CROPPING SYSTEMS" sections for specific recommended uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE, GREEN STEMS OR FRUIT OF DESIRABLE CROPS, PLANTS, TREES OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Except as otherwise specified on this label, repeat treatments must be made before the crop emerges in accordance with the instructions of this label.

Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts per acre of this product per year.

The maximum use rates stated throughout this product labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

For any crop not listed below, applications must be made at least 30 days prior to planting.

Do not harvest or feed treated vegetation for 8 weeks following application. Following spot treatment or selective equipment use, allow 14 days before grazing domestic livestock or harvesting forage grasses and legumes.

ROW CROPS

- CORN (ALL)*
- COTTON*
- PEANUTS
- SORGHUM (MILO)*
- SOYBEANS*
- SUGARCANE*

CEREAL GRAINS

- BARLEY*
- BUCKWHEAT*
- MILLET (PEARL, PROSO)*
- OATS*
- RICE**
- RYE*
- TRITICALE*
- WHEAT (ALL)*
- WILD RICE*

CITRUS

- CALAMONDIN
- CHIRONJA
- CITRON
- GRAPEFRUIT
- KUMQUAT
- LEMON
- LIME
- MANDARIN ORANGE
- ORANGE (ALL)
- PUMMELO
- TANGELO
- TANGERINE
- TANGORS

TREE NUTS

- ALMOND
- BEECHNUT
- BRAZIL NUT
- BUTTERNUT
- CASHEW
- CHESTNUT
- CHINQUAPIN
- FILBERT (HAZELNUT)
- HICKORY NUT
- MACADAMIA
- PECAN
- PISTACHIO
- WALNUT (BLACK, ENGLISH)

VINE CROPS

- GRAPES
- KIWI FRUIT

TREE FRUITS

- APPLE
- APRICOTS
- CHERRY (SWEET, SOUR)
- LOQUAT
- MAYHAW
- NECTARINE
- OLIVE
- PEACH
- PEAR
- PLUM/PRUNE (ALL)
- QUINCE

VEGETABLES

- ARTICHOKE,
- JERUSALEM
- ASPARAGUS*
- BEANS (ALL)
- BEET GREENS
- BEETS (RED,SUGAR)
- BROCCOLI (ALL)
- BRUSSELS SPROUTS
- CABBAGE (ALL)
- CABBAGE, CHINESE
- CANTALOUPE***
- CARROT
- CAULIFLOWER
- CASABA MELON***
- CELERIAC
- CELERY
- CHARD, SWISS
- CHICORY
- COLLARDS
- CRENSHAW MELON***
- CUCUMBER***
- EGGPLANT***
- ENDIVE
- GARLIC***
- GOURDS***
- GROUND CHERRY***
- HONEYDEW MELON***
- HONEY BALL MELON***
- HORSERADISH
- KALE
- KOHLRABI
- LEEK
- LENTILS
- LETTUCE
- MANGO MELON***
- MELONS (ALL)***
- MUSKMELON***
- MUSTARD GREENS
- OKRA
- ONION
- PARSLEY
- PARSNIPS
- PEAS (ALL)
- PEPPER (ALL)***
- PERSIAN MELON***
- POTATO (IRISH,SWEET)
- PUMPKIN***
- RADISH
- RAPE GREENS
- RHUBARB
- RUTABAGA
- SHALLOT
- SPINACH (ALL)
- SQUASH (SUMMER,WINTER)***
- TOMATILLO***
- TOMATO***†
- TURNIP
- WATERCRESS***
- WATERMELON***
- YAMS

SMALL FRUITS AND BERRIES

- BLACKBERRY
- BLUEBERRY
- BOYSENBERRY
- CRANBERRY
- CURRANT
- DEWBERRY
- ELDERBERRY
- GOOSEBERRY
- HUCKLEBERRY
- LOGANBERRY
- OLALLIEBERRY
- RASPBERRY (BLACK,RED)

FORAGE CROPS AND LEGUMES

ALFALFA*
FORAGE GRASSES*
FORAGE LEGUMES*

TROPICAL CROPS

ACEROLA
ATEMOYA
AVOCADO
BANANA (PLANTAINS)
BREADFRUIT
CANISTEL
CARAMBOLA
CHERIMOYA
COCOA BEANS
COFFEE
DATES
FIGS
GENIP
GUAVA
JABOTICABA
JACKFRUIT
LONGAN
LYCHEE
MANGO
PAPAYA
PASSION FRUIT
PERSIMMONS
PINEAPPLE****
POMEGRANATE
SAPODILLA
SAPOTE (BLACK, MAMEY, WHITE)
SOURSOP
SUGAR APPLE
TAMARIND
TEA

- * Spot treatments may be applied in these crops.
- ** Do not treat rice fields or levees when the fields contain flood water.
- *** Apply only prior to planting. Allow at least 3 days between application and planting.
- **** Do not feed or graze treated pineapple forage following application.
- † Use is restricted to direct seeded crops only.

When applying this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product, which could cause crop injury, from the plastic prior to transplanting. Residues can be removed by a single 0.5 inch application of water, either by natural rainfall or via a sprinkler irrigation system. Applications made at emergence will result in injury or death to emerged seedlings.

Spot Treatment (Only those crops with "*" can be spot treated.): Applications in growing crops must be made prior to heading of small grains and milo, initial pod set in soybeans, silking of corn, or boll opening on cotton.

For forage grasses and forage legumes see "SPOT TREATMENT" in the "PASTURES" section of "CROPPING SYSTEMS" in this label.

For dilution and rates of application using boom or hand-held equipment, see "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" and "WEEDS CONTROLLED" sections of this label.

NOTE: FOR FORAGE GRASSES AND FORAGE LEGUMES, NO MORE THAN ONE-TENTH OF ANY ACRE SHOULD BE TREATED AT ONE TIME. FOR ALL OTHER CROPS, DO NOT TREAT MORE THAN 10 PERCENT OF THE TOTAL FIELD AREA TO BE HARVESTED.

THE CROP RECEIVING SPRAY IN TREATED AREA WILL BE KILLED. TAKE CARE TO AVOID DRIFT OR SPRAY OUTSIDE TARGET AREA FOR THE SAME REASON.

Selective Equipment: This product may be applied through recirculating sprayers, shielded applicators or wiper applicators in cotton and soybeans. Shielded and wiper applicators may also be used in tree crops and grapes. Wiper applicators may be used in wheat, rutabagas, forage grasses and forage legumes, including pasture sites and grain sorghum (milo).

See the "SELECTIVE EQUIPMENT" part of the "APPLICATION EQUIPMENT and TECHNIQUES" section of this label for information on proper use and calibration of this equipment.

Allow at least the following time intervals between application and harvest:

Cotton, Soybeans	7 days
Apples, Citrus, Pear	1 day
Atemoya, Avocado, Breadfruit, Canistel, Carambola, Cherry, Grapes, Dates, Jaboticaba, Jackfruit, Longan, Lychee, Passion Fruit, Persimmons, Rutabagas, Sapodilla, Sapote, Soursop, Sugar Apple, Tamarind	14 days
Stone Fruit	17 days
Nut Crops	3 days
Wheat ¹	35 days
Sorghum (milo) ^{1,2}	40 days

¹ Do not use roller applicators.

² Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

ASPARAGUS

When applied as directed for "CROPPING SYSTEMS" under the conditions described, this product controls weeds listed on this label in asparagus.

For specific rates of applications and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

Prior to Crop Emergence: Apply this product prior to crop emergence for the control of emerged labeled annual and perennial weeds. **DO NOT APPLY WITHIN A WEEK BEFORE THE FIRST SPEARS EMERGE.**

Spot Treatment: Apply this product immediately after cutting, but prior to the emergence of new spears. Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

Postharvest: Apply this product after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears. Direct contact of the spray with the asparagus may result in serious crop injury.

NOTE: Select and use recommended types of spray equipment for postemergence postharvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of spray with the crop.

BERRIES AND SMALL FRUITS

Wiper applicators may be used in cranberries in accordance with instructions in this section.

For other berries, apply as a preplant broadcast application, or as a directed spray or wiper application post-planting.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for essential product performance information.

See the "SELECTIVE EQUIPMENT" part of the "APPLICATION EQUIPMENT and TECHNIQUES" section of this label for information on recommended use and calibration of this equipment.

Allow a minimum of 30 days between last application and harvest of cranberries. For other small fruits and berries, allow a minimum of 14 days between last application and harvest.

For Wick or other Wiper Applicators: Mix 1 gallon of this product in 4 gallons of water to prepare a 20 percent solution.

In severe infestations, reduce equipment ground speed to ensure that adequate amounts of this product are wiped on the weeds. A second treatment in the opposite direction may be beneficial.

Do not permit herbicide solution to contact desirable vegetation, including green shoots, canes or foliage.

CORN

Hooded Sprayers: This product may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

When applying to corn that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

Follow these requirements:

- The spray hoods must be operated on the ground or skimming across the ground.

- Do not apply more than 1 quart of this product per acre per application.
- Corn must be at least 12 inches tall, measured without extending leaves.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 mph.
- Maximum wind speed: 10 mph.
- Use low-drift nozzles.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weed Control Section" of this label.

Do not graze or feed corn forage or fodder following applications of this product through hooded sprayers.

Do not apply more than 3 quarts of this product per acre per year for hooded sprayer applications.

FALLOW AND REDUCED TILLAGE SYSTEMS

FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO SUPPLEMENTAL LABEL. Use this product in fallow and reduced tillage systems for control of annual weeds prior to emergence of crops listed in this label. Refer to the "WEEDS CONTROLLED" section of this label for specific rates and instructions. This product may be applied using ground or aerial spray equipment. See the "APPLICATION EQUIPMENT and TECHNIQUES" section of this label for instructions.

TANK MIXTURES

GLYPHOSATE plus "Banvel" plus Nonionic Surfactant

GLYPHOSATE plus 2,4-D plus Nonionic Surfactant

GLYPHOSATE plus "Goal" plus Nonionic Surfactant

DO NOT APPLY "BANVEL" OR 2,4-D TANK MIXTURES BY AIR IN CALIFORNIA.

Applications of 2,4-D or "Banvel" must be made at least 7 days prior to planting corn.

The addition of "Banvel" in a mixture with this product may provide short-term residual control of selected weed species. Some crop injury may occur if "Banvel" is applied within 45 days of planting. Refer to the "Banvel" and 2,4-D labels for cropping restrictions and other use instructions.

GLYPHOSATE plus "Goal" Tank Mixtures

This product alone or in tank mixtures with "Goal" plus 0.5 to 1 percent nonionic surfactant by total spray volume will provide control of those weeds listed below.

Make applications when weeds are actively growing and at the recommended stages of growth. Avoid spraying when weeds are subject to moisture stress, when dust is on the foliage or when straw canopy covers the weeds.

GLYPHOSATE 12 fluid oz/acre

Wheat	18"*
Barley	12"
Bluegrass, annual	6"
Barnyardgrass	6"
Rye	6"

GLYPHOSATE 16 fluid oz/acre

Annual grasses at left plus:

Ryegrass, annual	6"
Chickweed	6"
Groundsel	6"
Marestail	6"
Rocket, London	6"
Shepherdspurse	6"
Crabgrass	12"
Johnsongrass, seedling	12"
Lambsquarters	12"
Oats, wild	12"
Pigweed, redroot	12"
Mustards	12"

NOTE: Use 32 fluid ounces of this product per acre where heavy weed densities exist.

GLYPHOSATE	GLYPHOSATE
12 fluid oz/acre	16 fluid oz/acre
+	+
"Goal"***	"Goal"***
2 to 4 fluid oz/acre	2 to 4 fluid oz/acre

Annual grasses above plus:	Annual weeds above plus:
Cheeseweed, common 3"	Cheeseweed, common 6"
Chickweed 3"	Groundsel 6"
Groundsel 3"	Chickweed 12"
Rocket, London 6"	Rocket, London 12"
Shepherdspurse 6"	Shepherdspurse 12"

NOTE: Use 32 fluid ounces of this product per acre in mixtures with 2 to 4 fluid ounces of "Goal" per acre where heavy weed densities exist.

* Maximum height or length in inches.

** Use the higher rate of "Goal" when weeds approach maximum recommended height or stands are dense.

These recommended tank mixtures may be applied using ground or aerial spray equipment. Refer to the "WEEDS CONTROLLED" section of this label for specific rates and instructions.

ECOFARMING SYSTEMS

The recommendations made in this section are not registered for use in California.

The Ecofarming System consists of the following rotation: winter wheat, corn/sorghum, ecofallow.

Use the following tank mixtures for control of emerged annual weeds before planting corn or sorghum in the Ecofarming System.

GLYPHOSATE at 16 to 20 fluid ounces per acre

plus

2,4-D at 0.375 to 0.5 pound a.i. per acre

plus

Atrazine at 0.75 to 1 pound a.i. per acre

plus

"Lasso" at 2.5 to 3 quarts per acre

The above tank mixture should be applied in 28-0-0 or 32-0-0 liquid fertilizer carrier at 20 to 30 gallons per acre. The liquid fertilizer may be diluted with water to achieve the desired carrier volume.

WEEDS CONTROLLED - The following weeds, up to a maximum height of 4 inches, will be controlled:

Brome, downy,	<i>Bromus tectorum</i>
Cheat,	<i>Bromus secalinus</i>
Foxtail, green,	<i>Setaria viridis</i>
Foxtail, yellow,	<i>Setaria lutescens</i>
Kochia*,	<i>Kochia scoparia</i>
Lettuce, prickly,	<i>Lactuca serriola</i>
Pigweed, redroot,	<i>Amaranthus retroflexus</i>
Thistle, Russian,	<i>Salsola kali</i>
Wheat, volunteer,	<i>Triticum aestivum</i>

*For improved control of kochia, add 4 fluid ounces per acre (0.125 pound a.i. per acre) of "Banvel" to the above tank mixture.

Risk of crop injury from 2,4-D or "Banvel" can be reduced by applying this treatment 7 to 14 days before planting.

Refer to the label booklet for "Lasso" herbicide for preemergence weed control achieved by this tank mixture.

Refer to the specific product labels for crop rotation restrictions and cautionary statements for all products used in these tank mixtures.

AID TO TILLAGE

This product, when used in conjunction with preplant tillage practices, will provide control of downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 8 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Make applications when weeds are actively growing and before they are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage. Tank mixtures with residual herbicides may result in reduced performance.

POSTHARVEST GRAIN SORGHUM, SORGHUM REGROWTH CONTROL

This product may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 quart of this product per acre for control, or 1.5 pints of this product per acre for suppression. Use 0.5 percent nonionic surfactant in 3 to 10 gallons of spray solution per acre.

PASTURES

Apply this product prior to planting forage grasses and legumes.

Pasture or Hay Crop Renovation - When applied as a broadcast spray, this product controls the annual and perennial weeds listed in this label prior to planting forage grasses or legumes. Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Spot Treatment - When applied as a spot treatment as recommended, this product controls annual and perennial weeds listed in this label which are growing in pastures, forage grasses and forage legumes composed of bahiagrass, bermudagrass, bluegrass, brome, fescue, orchardgrass, ryegrass, timothy, wheatgrass, alfalfa or clover.

Wiper Application - when applied as directed, this product controls or suppresses the weeds listed under "WIPER APPLICATORS" in the "SELECTIVE EQUIPMENT" section of this label.

For spot treatment and wiper application, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre should be treated at one time. Further applications may be made in the same area at 30-day intervals. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

SUGARCANE

When applied as directed for "CROPPING SYSTEMS", under the conditions described, this product controls those emerged annual and perennial weeds listed on this label growing in or around sugarcane or in fields prior to the emergence of plant cane. This product will also control undesirable sugarcane.

NOTE: Where repeat treatments are necessary, do not exceed a total of 10.6 quarts of this product per acre per year. Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

Broadcast Treatment - Apply this product in 10 to 40 gallons of water per acre on emerged weeds prior to the emergence of plant cane.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

For removal of last stubble or ratoon cane, apply 4 to 5 quarts of this product in 10 to 40 gallons of water per acre to new growth having at least 7 or more new leaves. Allow 7 or more days after application before tillage.

Spot Treatment in or Around Sugarcane Fields - For dilution and rates of application using hand-held equipment, see "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" and "WEEDS CONTROLLED" sections of this label.

For control of volunteer or diseased sugarcane, make a 1 percent solution of this product in water and spray to wet the foliage of vegetation to be controlled.

NOTE: When spraying volunteer or diseased sugarcane, the plants should have at least 7 new leaves.

Avoid spray contact with healthy cane plants since severe damage or destruction may result.

Do not feed or graze treated sugarcane forage following application.

CONSERVATION TILLAGE, MINIMUM TILLAGE AND NO-TILL SYSTEMS

CORN AND SOYBEANS TANK MIXTURES

The recommendations made in this section are not registered for use in California.

When applied as recommended under the conditions described, the tank mixtures listed in this section control many emerged weeds, and give preemergence control of many annual weeds where corn or soybeans will be planted directly into a cover crop, established sod or in previous crop residues.

Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures. For mixing instructions, see the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label.

Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre before, during or after planting. Do not apply these mixtures after crop emergence.

When tank mixing with residual herbicides, add an agriculturally approved nonionic surfactant at 0.5 to 1 percent by volume of spray solution. The addition of 1 to 2 percent dry ammonium sulfate by weight may increase the performance of this product.

NOTE: When using these tank mixtures, do not exceed 4 quarts of this product per acre.

CORN

For residual control, this product may be tankmixed with the following herbicides or combination of herbicides:

"Lasso"/Alachlor	Atrazine
"Lariat"	Cyanazine
"Bullet"	"Simazine"
"Dual"	"Prowl"
"Bicep"	"Micro-Tech"
"Partner" ¹	

For improved burndown, this product may be tankmixed with 2,4-D or dicamba. Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn. See the "WEEDS CONTROLLED" section for specific rate information.

¹ "Partner" herbicide is not registered in California.

SOYBEANS

For residual control, this product may be tankmixed with the following herbicides or combination of herbicides:

CANOPY®	“Lorox Plus”
“Command”	“Preview”
“Dual”	“Prowl”
“Gemini”	“Turbo”
“Lasso”/Alachlor	“Scepter”
LEXONE®	“Sencor”
Linuron	“Squadron”
“Pursuit”	“Pursuit Plus”
“Partner”	“Micro-Tech”

For improved burndown, this product may be tankmixed with the following herbicides:

2,4-DB

2,4-D*

* See the label for 2,4-D for intervals between application and planting.

CORN AND SOYBEANS

Annual Weeds--For difficult-to-control weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1 to 1.5 pints of this product per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall. For a complete list of annual weeds controlled, see the "WEEDS CONTROLLED" section of this label.

Perennial Weeds--At normal application times in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. See the "WEEDS CONTROLLED" section of this label for the proper stage of growth for perennial weeds.

Use of 2 to 4 quarts of this product per acre in the tank mixtures mentioned above, under these conditions provides top kill and reduces competition from many emerged perennial grass and broadleaf weeds. For emerged perennial weeds controlled, see the "WEEDS CONTROLLED" section of this label.

To obtain the desired stage of growth, it may be necessary to apply this product alone in the late summer or fall and then follow with a label-approved, seedling weed-control program at planting.

USE OF THESE TANK MIXTURES FOR BERMUDAGRASS OR JOHNSONGRASS CONTROL IN MINIMUM TILLAGE SYSTEMS IS NOT RECOMMENDED. For bermudagrass control, follow the instructions under "CONTROL OF PERENNIAL WEEDS" section of this label and then use a label-approved, seedling weed-control program in a minimum tillage or conventional tillage system. For Johnsongrass control, follow instructions under "CONTROL OF PERENNIAL WEEDS" section of this label, and then use a label-approved, seedling weed-control program with conventional tillage.

PREHARVEST APPLICATIONS

When applied as directed under the conditions described, this product controls annual and perennial weeds listed on this label prior to the harvest of cotton, soybeans and wheat.

For specific rates and application instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

This product may be applied by both ground and aerial application equipment. **DO NOT APPLY MORE THAN 1 QUART PER ACRE OF THIS PRODUCT BY AIR.** See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for instructions for ground and aerial applications.

NOTE: Do not apply to crops grown for seed. Reduction in germination or vigor may occur.

The use of this product for preharvest grain sorghum (Milo) is not registered in California.

SOYBEANS

Apply after pods have set and lost all green color. Allow a minimum of 7 days between application and harvest of soybeans. Care should be taken to avoid excessive seed shatter loss due to ground application equipment.

Do not graze or harvest treated crop for livestock feed within 25 days of last preharvest application.

DO NOT APPLY MORE THAN 6 QUARTS PER ACRE OF THIS PRODUCT FOR PREHARVEST APPLICATIONS.

COTTON

Broadcast Applications--This product may be applied using either aerial or ground spray equipment. For ground applications with broadcast equipment, apply this product in 10 to 20 gallons of water per acre. For aerial applications, apply this product in 3 to 10 gallons of water per acre.

This product provides weed control and cotton regrowth inhibition when applied prior to the harvest of cotton. Apply 1 to 2 quarts of this product in 3 to 10 gallons of water per acre for cotton regrowth inhibition. Do not apply more than 2 quarts of this product per acre for preharvest applications. **THE USE OF ADDITIVES FOR PREHARVEST APPLICATION TO COTTON IS PROHIBITED.**

This product may be tank mixed with "DEF 6", "Folex", or "Prep" to provide additional enhancement of cotton leaf drop.

Allow a minimum of 7 days between application and harvest of cotton.

Apply after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.

Do not feed or graze treated cotton forage or hay following preharvest applications.

GRAIN SORGHUM (MILO)

Make applications at 30% grain moisture or less and at least 7 days prior to harvest.

Apply up to 2 quarts of this product per acre.

WHEAT

Apply after the hard-dough stage of grain (30% or less grain moisture) and at least 7 days prior to harvest.

DO NOT APPLY MORE THAN 1 QUART PER ACRE OF THIS PRODUCT FOR PREHARVEST APPLICATIONS TO WHEAT.

TREE AND VINE CROPS

This product is recommended for weed control in established groves, vineyards, and orchards, or for site preparation prior to transplanting crops listed in this section. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed in this section. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for specific information on use of equipment.

When applying this product, refer to the "WEEDS CONTROLLED" section of this label and to specific recommendations in this section for rates to be used.

NOTE: Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. This product does not provide residual weed control. For subsequent weed control, use repeated applications of this product. Do not apply more than 10.6 quarts of this product per acre per year.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

AVOID PAINTING OUT STUMPS WITH THIS PRODUCT AS INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut and have not been allowed to regrow to the recommended stage for treatment.

For specific rates of applications and instructions, see the "WEEDS CONTROLLED" section of this label, and to specific recommendations which follow.

MIDDLES MANAGEMENT

FOR ANNUAL WEEDS IN MIDDLES BETWEEN ROWS OF TREE AND VINE CROPS

For citrus crops, treat uniformly between trees.

GLYPHOSATE

GLYPHOSATE plus "Goal"

This product alone or in mixtures with "Goal" will control or suppress the annual weeds listed below.

Apply the recommended rates of this product, either alone or in mixtures with "Goal", plus 0.5 to 1 percent nonionic surfactant by spray volume in 3 to 10 gallons of water per acre. Apply when weeds are actively growing and less than 6 inches in height or diameter. If weeds are under drought stress, irrigate prior to application. Reduced control may occur if weeds have been mowed prior to application. Up to 48 fluid ounces per acre of this product may be used to control weeds, which have been mowed, are stressed or are growing in dense populations.

WEED SPECIES	MAXIMUM HEIGHT/DIA. (INCHES)	RATE PER ACRE	
		GLYPHOSATE (Fl Oz)	"Goal" (Fl Oz)
Barley <i>Hordeum vulgare</i>	6	8	--
Bluegrass, annual <i>Poa annua</i>			
Barnyardgrass <i>Echinochloa crus-galli</i>	6	12	--
Chickweed, common <i>Stellaria media</i>			
Red Maids <i>Calandrinia ciliata</i>			
Crabgrass <i>Digitaria spp.</i>	6	16	--
Fleabane, hairy <i>Conyza bonariensis</i>			
Groundsel, common <i>Senecio vulgaris</i>		16 to 32	+ 4 to 16**
Junglerice <i>Echinochloa colonum</i>			
Lambsquarters, common <i>Chenopodium album</i>			
Pigweed, redroot <i>Amaranthus retroflexus</i>			
Rocket, London <i>Sisymbrium irio</i>			
Ryegrass, common <i>Lolium multiflorum</i>			
Shepherdspurse <i>Capsella bursa-pastoris</i>			
Sowthistle, annual <i>Sonchus oleraceus</i>			
Cheeseweed, common <i>Malva spp</i>	3	12 to 32	+ 4 to 16
Cheeseweed, common <i>Malva spp</i>	6	16 to 32	+ 4 to 16
Filaree* <i>Erodium spp.</i>			
Horseweed/Marestail <i>Conyza canadensis</i>			
Nettle, stinging <i>Urtica dioica</i>			
Purselane, common* <i>Purtulaca oleracea</i>			

* Suppression only.

** The mixture of this product plus "Goal" is recommended when weeds are stressed or growing in dense populations.

STRIPS

FOR ANNUAL AND PERENNIAL WEEDS IN STRIPS OF TREE AND VINE CROPS

TANK MIXTURES WITH RESIDUAL HERBICIDES

When applied as a tank mixture, this product provides control of the emerged annual weeds and control or suppression of emerged perennial weeds listed in this label. The following residual herbicides will provide preemergence control of those weeds listed in the individual product labels.

GLYPHOSATE plus:

“Goal 2XL”

“Karmex DF”

KROVAR® I

KROVAR® II

“Simazine”

“Princep Caliber 90”

“Simazine 4L”

“Simazine 80W”

“Solicam 80DF”

“Surflan AS”

“Surflan 75W”

“Simazine” (80W, or 4L, or “Princep Caliber 90”) plus

“Surflan” (AS or 75W)

“Goal 2XL” plus “Surflan” (AS or 75W)

“Goal 2XL” plus “Simazine” (80W, or 4L, or “Princep Caliber 90”)

“Goal 2XL” plus “Surflan” (AS or 75W) plus “Simazine” (80W, 4L, or “Princep Caliber 90”)

Do not apply these tank mixtures in Puerto Rico.

When tankmixing with residual herbicides, add an agriculturally approved nonionic surfactant at 0.5 to 1 percent by volume of spray solution.

Refer to the individual product labels for specific crops, rates, geographical restrictions and precautionary statements.

Read and carefully observe the label claims, cautionary statements, rates and all other information on the labels of all products.

RECOMMENDED RATES

Annual Weeds - Apply 1 to 5 quarts per acre of this product in these tank mixtures. Use rates at the higher end of the recommended range when weeds are stressed, growing in dense populations or are greater than 12 inches tall.

Perennial Weeds - Apply 1 pint to 5 quarts per acre of this product in these tank mixtures to control or suppress perennial weeds. Follow the recommendations in the "WEEDS CONTROLLED" section of this label for stage of growth and application rates for specific perennial weeds.

GLYPHOSATE plus “Goal” plus “Simazine”/“Surflan”

This product plus low rates of “Goal” in 3-way or 4-way mixtures with “Simazine” and/or “Surflan” will provide postemergence control of the weeds listed below.

Refer to the individual “Simazine” and “Surflan” labels for preemergence rates, weeds controlled, precautionary statements and other important information.

Apply these tank mixtures in 3 to 40 gallons of water. Add 0.5 to 1 percent nonionic surfactant by total spray volume to the spray solution.

Apply 1 to 5 quarts per acre of this product plus 4 to 48 fluid ounces per acre of “Goal” plus labeled rates of “Simazine” and/or “Surflan” to control the following weeds:

Barley, wild,	<i>Hordeum leporinum</i>
Bluegrass, annual,	<i>Poa annua</i>
Cheeseweed, common,	<i>Malva spp.</i>
Chickweed, common,	<i>Stellaria media</i>
Filaree*,	<i>Erodium spp.</i>
Fleabane, hairy,	<i>Conyza bonariensis</i>
Groundsel, common,	<i>Senecio vulgaris</i>
Horseweed/Marestail,	<i>Conyza canadensis</i>
Nettle, stinging,	<i>Urtica dioica</i>
Pineappleweed,	<i>Matricaria matricarioides</i>
Rocket, London,	<i>Sisymbrium irio</i>
Shepherdspurse,	<i>Capsella bursa-pastoris</i>
Sowthistle, annual,	<i>Sonchus oleraceus</i>

* Use a minimum of 1.5 quarts of this product in these mixtures.

NOTE: This recommendation does not preclude the use of “Goal” in these mixtures at higher, labeled rates for preemergence weed control.

PERENNIAL GRASS SUPPRESSION ORCHARD FLOORS

When applied as directed, this product will suppress vegetative growth as indicated below.

Bahiagrass

This product will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with a single application and approximately 120 days with sequential applications. Apply this product 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 6 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 25 gallons of water per acre.

Sequential applications of this product plus nonionic surfactant may be made at approximately 45-day intervals to extend the period of seedhead and vegetative growth suppression. For continued seedhead suppression, sequential applications must be made prior to seedhead emergence. Apply no more than 2 sequential applications per year. As a first sequential application, apply 4 fluid ounces of this product plus nonionic surfactant. A second sequential application of 2 to 4 fluid ounces may be made approximately 45 days after the last application.

Bermudagrass

For burndown, apply 1 to 2 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 20 gallons of water per acre. Use 1 quart of this product in 3 to 20 gallons of water per acre east of the Rocky Mountains. Use 1 to 2 quarts of this product in 3 to 10 gallons of water per acre west of the Rocky Mountains. Use this treatment only if reduction of the bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

Suppression only (east of the Rocky Mountains): Apply 6 to 16 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 20 gallons of water per acre no sooner than 1 to 2 weeks after full green-up. Mowing prior to application may occur provided a minimum height of 3 inches is maintained. Rates of 6 to 10 fluid ounces of this product plus nonionic surfactant should be used in shaded conditions or where a lesser degree of suppression is desired. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated.

Suppression only (west of the Rocky Mountains): Apply 16 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre to bermudagrass up to 6 inches in height and no sooner than 1 to 2 weeks after full green-up. Mowing prior to application may occur provided a minimum height of 3 inches is maintained. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated.

Cool Season Grass Covers

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 20 gallons of water per acre. For best suppression, add ammonium sulfate to the spray solution at a rate of 2 percent by weight or 17 pounds per 100 gallons of spray solution.

For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant. Do not add ammonium sulfate.

For best results, mow cool-season grass covers in the spring to even their height and apply the recommended rate of this product 3 to 4 days after mowing. Avoid treating cool season grass covers under poor growing conditions, such as drought stress (drip irrigation), disease or insect damage.

LOW VOLUME APPLICATION (FLORIDA AND TEXAS)

For burndown or control of the weeds listed, apply the recommended rates of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

ANNUAL WEEDS

Goatweed--Apply 2 to 3 quarts per acre of this product plus 17 pounds of ammonium sulfate per 100 gallons of water plus 0.5 to 1 percent nonionic surfactant by total spray volume. Apply in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are less than 8 inches tall and 3 quarts per acre when plants are greater than 8 inches. If goatweed is greater than 8 inches tall, the addition of KROVAR® II or "Karmex" may improve control. Use labeled rates for these residual products.

Read and carefully observe the label claims, cautionary statements, rates and all other information on the KROVAR® II and "Karmex" labels.

PERENNIAL WEEDS

Apply when weeds are actively growing and at the growth stages listed in the "PERENNIAL WEEDS CONTROLLED" section of this label. If perennial weeds are mowed, allow weeds to regrow to the recommended stage of growth.

S = Suppression

B = Burndown

PC = Partial control

C = Control

WEED SPECIES	GLYPHOSATE RATE PER ACRE			
	1 qt	2 qts	3 qts	5 qts
Bermudagrass	B	•	PC	C
Guineagrass				
Texas and Florida Ridge Florida	B	C	C	C
Flatwoods	•	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	•	PC	C

TREE CROPS

Citrus*****: calamondin, chironja, citron, grapefruit, kumquat, lemon, lime, mandarin orange, orange, pummelo, tangelo, tangerine, tangors.

Nuts***: almond, beechnut, Brazil nut, butternut, cashew, chestnuts, chinquapin, filbert, hazel nut, hickory nut, macadamia, pecan, pistachio, walnut.

Pome Fruit*****: apple, loquat, mayhaw, pear, quince.

Stone Fruit*****: apricots, cherries, nectarines, olives, peaches, plums/prunes.

For cherries, any application equipment listed in this section may be used in all states.

For citron and olives, apply as a directed spray only.

Any application equipment listed in this section may be used in apricots, nectarines, peaches and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah and Washington, except for peaches grown in the states specified in the following paragraph. In all other states use wiper equipment only.

For PEACHES grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee only, apply with a shielded boom sprayer or shielded wiper applicator, which prevents any contact of this product with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees which have been planted in the orchard for 2 or more years. **EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED.**

Tropical Fruit: acerola*, atemoya*, avocado*, banana (plantains)****, breadfruit*, canistel*, carambola*, cherimoya*, cocoa beans*, coffee****, dates*, figs*, genip*, guava****, jaboticaba*, jackfruit*, longan*, lychee*, mango*, mayhaw*, papaya****, passion fruit*, persimmons*, pomegranate*, sapodilla*, sapote*, soursop*, sugar apple*, tamarind*, tea*. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

NOTE:

- * Allow a minimum of 14 days between last application and harvest.
- ** Allow a minimum of 3 days between last application and harvest.
- *** Allow a minimum of 17 days between last application and harvest.
- **** Allow a minimum of 28 days between last application and harvest.
- ***** Allow a minimum of 1 day between last application and harvest.

VINE CROPS

Kiwi Fruit

Grapes: Any variety of table, wine or raisin grape may be treated with any equipment listed in this section.

Applications should not be made when green shoots, canes, or foliage are in the spray zone.

Allow a minimum of 14 days between last application and harvest.

In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.

FOR COTTON WITH THE “ROUNDUP READY”, GENE IN-CROP APPLICATIONS

General Information

DUPONT RECOMMENDS THIS PRODUCT FOR USE ONLY OVER-THE-TOP OF OR DIRECTED ONTO IMPROVED COTTON VARIETIES THAT ARE DESIGNATED AS COTTON WITH THE “ROUNDUP READY” GENE. SEVERE INJURY OR DEATH OF COTTON WILL RESULT IF ANY COTTON VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE “ROUNDUP READY” GENE ARE SPRAYED WITH THIS PRODUCT. AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR ANY DESIRABLE PLANTS AND TREES OTHER THAN CROPS WITH THE “ROUNDUP READY” GENE, SINCE SEVERE INJURY OR DESTRUCTIONS WILL RESULT.

USE OF GLYPHOSATE IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF “ROUNDUP READY” COTTON, HOWEVER, VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS MAKE IT IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

“Roundup Ready” cotton varieties must be purchased from an authorized licensed seed supplier. The designation, “Roundup Ready”, indicates the cotton variety contains a patented proprietary trait.

For over-the-top uses on “Roundup Ready” crop varieties, crop safety and weed control performance are not warranted by DuPont when this product is used in conjunction with “Brown Bag” or “Bin-Run” seed saved from previous year’s production and replanted.

Application Instructions

This product will control many troublesome weeds with over-the-top, post-directed, hooded sprayer, or preharvest applications in Roundup Ready cotton.

Maximum Allowable Yearly Rates

1. Combined total per year for all applications: 8 quarts/A
2. Preplant, Preemergence applications: 5 quarts/A
3. Total in-crop applications from cracking to layby: 4 quarts/A
4. Maximum preharvest application rate: 2 quarts/A

For ground applications with broadcast equipment, apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

For aerial applications, apply this product in 3 to 15 gallons of water per acre.

Do not exceed a maximum rate of 1 quart per acre of this product when making applications by air, unless otherwise directed.

Extreme care must be used when applying this product to prevent injury to desirable plants and crops which do not contain the "Roundup Ready" gene. See Spray Drift Management section of this label for instructions.

There are no rotational crop restrictions following applications of this product.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product to "Roundup Ready" cotton. Follow the cleaning procedures specified on the label of the product(s) previously used. Cotton is very sensitive to many herbicides at extremely low concentrations and care should be taken to thoroughly clean all equipment prior to use.

Over-the-top applications: This product may be applied by aerial or ground application equipment postemergence to "Roundup Ready" cotton from the ground cracking stage until the four leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). Over-the-top applications made after the four leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss. Any single over-the-top broadcast application should not exceed 1 quart per acre. No more than two over-the-top broadcast applications may be made from crop emergence through the four leaf (node) stage of development. Sequential over-the-top applications of this product must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

NOTE: Always plant into a weed free seedbed. In no-till and stale seedbed systems, always burn down existing weeds before cotton emerges. Apply a preplant burndown treatment of 16-48 fluid ounces per acre of this product.

Post-directed or hooded applications: This product may be applied using precision post-directed or hooded sprayers to "Roundup Ready" cotton through layby. At this stage, post-directed equipment should be used which directs the spray to the base of the cotton plants. Contact of the spray with cotton leaves should be avoided to the maximum extent possible. To minimize spray onto the leaves of the cotton plants, place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row, and maintain low spray pressure (less than 30 PSI). For best results, make applications while weeds are small (less than 3 inches). Any single post-directed application should not exceed 1 quart per acre of this product. No more than two applications should be made from the fifth leaf through layby. Sequential in-crop applications of this product must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

Salvage Treatment: This treatment may be used after the four leaf stage of development and should only be used where weeds threaten to cause the loss of the crop. One quart per acre may be applied either as an over-the-top application or as a post-directed treatment sprayed higher on the cotton plants and over the weeds. **NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT SHOULD BE USED PER GROWING SEASON.**

Preharvest applications: This product may be applied for preharvest annual and perennial weed control as a broadcast treatment to "Roundup Ready" cotton after 20% boll crack. For application rates please see the "WEEDS CONTROLLED" section of the GLYPHOSATE label booklet. This product may be applied using either aerial or ground spray equipment. Aerial and ground applications may be made up to a maximum of 2 quarts per acre. Allow a minimum of 7 days between final application and harvest of cotton or feeding of cotton forage or hay. **THE USE OF ADDITIVES FOR PREHARVEST APPLICATION OF GLYPHOSATE TO "ROUNDUP READY" COTTON IS PROHIBITED. NOTE: GLYPHOSATE will not enhance the performance of harvest aids when applied to "Roundup Ready" cotton. DO NOT APPLY GLYPHOSATE PREHARVEST TO CROPS GROWN FOR SEED.**

Weeds controlled: For specific rates of application and instructions for control of various annual and perennial weeds, refer to "WEEDS CONTROLLED" section of this label. GLYPHOSATE applied at 1 quart per acre will burndown or suppress the growth of the following perennial weeds and reduce crop competition: yellow and purple nutsedge, rhizome johnsongrass, common bermudagrass, silverleaf nightshade, trumpet creeper, and redvine. Fall preharvest applications may be required for control of these perennial weeds.

TANK MIXTURES

GLYPHOSATE plus DuPont STAPLE® (GLYPHOSATE Tolerant Cotton)

A tank mixture of STAPLE® at 0.6 ounces per acre plus GLYPHOSATE at 24 ounces per acre may be applied as an early postemergence treatment in glyphosate tolerant cotton, such as “Roundup Ready” cotton, for improved control of hemp sesbania, morningglory (entireleaf, ivyleaf, pitted, scarlet/red), prickly sida and palmer pigweed. The addition of STAPLE® to a “Roundup Ready” program will provide residual control of prickly sida, pigweed, spotted spurge, velvetleaf and spurred anoda. Rainfall (0.5-1 inch) following the postemergence application is required for residual control.

GLYPHOSATE plus STAPLE® may be applied postemergence to glyphosate tolerant cotton, such as “Roundup Ready” cotton, through the 4th true leaf stage of growth (when 5th true leaf is the size of a quarter or less).

Refer to the label for further application information and use restrictions. Follow the label guidelines that are the most restrictive.

NOTE: No antagonism has been observed to annual grass species from this tank mixture.

Some weeds with multiple germination times or suppressed (stunted) weeds may require sequential applications of this product for control.

Tank mixtures with other herbicides not recommended on this label may result in reduced weed control or crop injury and are not recommended for over-the-top applications of this product.

SPRAYER PREPARATION

It is important that spray equipment is clean and free of existing pesticide deposits before using GLYPHOSATE. Follow the cleanup procedures specified on the label of the product(s) previously used. If no cleanout procedure is provided, follow this cleanout procedure for all application equipment before using GLYPHOSATE:

1. Thoroughly rinse sprayer, tanks, boom and hoses with clean water.
2. Partially fill tank with water and add ammonia (1 gal. of ammonia per 100 gal. of tank volume) or a tank cleaner. Complete filling the tank and flush the cleaning solution through the boom hoses. Let solution stand for 15 minutes while agitating/recirculating, and then drain the tank by flushing the hoses, booms and nozzles.
3. Thoroughly rinse the sprayer, tanks, boom and hoses with clean water.
4. Follow label directions on product(s) previously sprayed for disposal.

Remove cap from container. Remove water soluble powder packet and place into the necessary volume of water (see Spray Equipment-Volumes) in the spray tank with the agitator running. Do not attempt to open or divide the contents of the water soluble packet. Do not tip or invert

the bottle to remove the water soluble packet. Doing so could dump the insert into the spray tank and spill the liquid. Remove insert from bottle, rinse thoroughly and discard. Pour the entire contents of the bottle containing the soluble liquid into spray tank. Continuous agitation is required for a uniform suspension and application.

PRECAUTION: Do not use chlorine bleach with ammonia. See Sprayer Cleanup Section for more information.

SPRAYER CLEANUP

Spray equipment must be clean and free of previous pesticide deposits before applying. Using the cleanup procedures specified on the label of the previously used product, clean all application equipment before applying. If no cleanup procedure is provided, use the procedure that follows. Immediately following applications thoroughly clean all mixing and spray equipment according to the following instructions:

1. Drain Tank: Thoroughly hose down the interior surfaces of the tank; then flush tank, boom and hoses with clean water for a minimum of 5 minutes. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and add one gal. of household ammonia* (3% active) for every 100 gal. of water. Flush the cleaning solution through the boom, hoses and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 minutes. Again, flush the boom, hoses and nozzles with the cleaning solution, then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing the cleaning agent and water.
4. Repeat step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing the water through the hoses and boom.
6. Dispose of the rinsate on site or at an approved waste disposal facility.

* Equivalent amounts of an alternate-strength ammonia solution or DuPont approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instruction.

PRECAUTION

Do not use chlorine bleach with ammonia when cleaning out spray tanks. All traces of liquid fertilizer containing ammonia, ammonia nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding any chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odor which can cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (greater than 150–200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions! See the Wind, Temperature and Humidity, and Temperature Inversions sections below.

CONTROLLING DROPLET SIZE

GENERAL TECHNIQUES

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use a higher-capacity nozzle instead of increasing pressure.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

AIRCRAFT

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.

BOOM LENGTH AND HEIGHT

- **Boom Length (aircraft)** - The boom length should not exceed 3/4 of the wing length, using shorter booms decreases drift potential. For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.
- **Boom Height (aircraft)** - Application more than 10 ft above the canopy increases the potential for spray drift.
- **Boom Height (ground)** Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they effect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

SURFACE TEMPERATURE INVERSIONS

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift, and not interfering with uniform deposition of the product.

FOR AERIAL APPLICATION IN ARKANSAS ONLY

Use the recommended rate of this product in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range are recommended.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75% of the length of the wingspan or rotor. In many cases, reducing this distance to 65% of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing air flow on rotary winged aircraft. Avoid the use of nozzles with wide angle discharge.

Do not apply this product when winds are in excess of 10 miles per hour.

Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 mph.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

FOR AERIAL APPLICATION IN CALIFORNIA ONLY

Aerial applications of this product are allowed in the following situations:

1. In fallow and reduced tillage systems prior to the emergence or transplanting of labeled crops.
2. In cotton, prior to harvest. Refer to the cotton instructions section for specific preharvest application instructions.

Do not plant subsequent crops other than those listed on this label for this product for 30 days following application.

When applied as recommended, under the conditions described, this product controls annual and perennial weeds listed in the label booklet.

DO NOT EXCEED A MAXIMUM RATE OF 2 QUARTS PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR IN FALLOW AND REDUCED TILLAGE SYSTEMS.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR IN COTTON, PRIOR TO HARVEST.

AERIAL EQUIPMENT

Use the recommended rates of this product in 3 to 15 gallons of water per acre.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation may require buffer zones in excess of 500 feet.

4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the precautionary statements and all other information appearing on the additive label.

Ensure uniform application—to avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills, **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

FOR AERIAL APPLICATION IN FRESNO COUNTY, CA ONLY

FROM FEBRUARY 15 THROUGH MARCH 31 ONLY

For aerial application outside these dates, refer to the "FOR AERIAL APPLICATION IN CALIFORNIA ONLY" section of this label.

Applicable Area

These instructions only apply to the area contained inside the following boundaries within Fresno County, California only.

North:Fresno County line
South:Fresno County line
East:State Highway 99
West:Fresno County line

General Information

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of GLYPHOSATE herbicide. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor, and aerial applicator.

Written Recommendations

A written recommendation **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation **MUST** state the proximity of surrounding crops, and that conditions of each manufacturer's applicable product label(s) and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of GLYPHOSATE herbicide is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight, and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved "fly-ins" constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

Application at night - Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal. Keep container closed to prevent spills and contamination.

PRODUCT DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.

Bulk Containers: Triple rinse emptied bulk containers. Then offer for recycling or reconditioning, or dispose of in a manner approved by state and local authorities.

Plastic 1-Way Containers: Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

Registered trademark of:

"Bullet", "Lasso", "Micro-Tech", "Partner", and "Lariat", "Roundup" are trademarks of Monsanto Company.

"Lorox Plus", and "Karmex" are trademarks of Griffin LLC

"Dual", "Bicep", "Princep Caliber 90", "Solicam" are trademarks of Novartis Corporation.

"Surflan" is a trademark of DowElanco Products Company.

"Banvel" is a trademark of BASF, Inc.

"Folex" and "Prep" are trademarks of Rhone-Poulenc, Inc.

"Goal" is a trademark of Rohm and Haas Company.

"Sencor" and "Turbo" are trademarks of Bayer AG.

"Prowl", "Pursuit", "Pursuit Plus", "Scepter" and "Squadron" are trademarks of American Cyanamid Company.

"Command" is a trademark of FMC Corporation.

"DEF 6" is a trademark of Mobay Chemical Company.

"Simazine" is a registered product of United Agri Products.

SL - 746A 061400 07-06-00

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks.
WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: www.dupont.com/ag/us

© 2000 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898