

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Stinger*

EPA Reg. No. 62719-73

EPA 24(c) Special Local Need Registration SLN MT-980006 For Distribution and Use Only in the State of Montana

Revised Preharvest Interval for Sugar Beets

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Stinger before applying. Carefully follow all precautionary statements and applicable use directions.
- In addition to restrictions and limitations imposed by this supplemental labeling, use of Stinger is subject to all use precautions and limitations imposed by the label affixed to the container for Stinger.

Directions for Use

Stinger* herbicide is recommended for the control of various annual and perennial broadleaf weeds infesting sugar beets. Apply 1/4 to 2/3 pint of Stinger per acre as a broadcast foliar spray. Apply when the sugar beets are in the cotyledon to 8 leaf stage of growth and the weeds are young and actively growing. Re-treat as necessary, but do not exceed 2/3 pints of Stinger per acre per season.

Do not apply within 90 days before harvest of beet roots and tops.

Follow directions in main product label for ideal rates and timing for control of specific weeds.

Use Precautions:

- **Chemigation:** Do not apply this product through any type of irrigation system.
- Avoid spray drift since small quantities of the spray may injure susceptible broadleaf crops. Refer to the main product label for recommendations on reducing spray drift.
- Carefully follow rotational crop restrictions and other use precautions and limitations on the product label for Stinger.

*Trademark of Dow AgroSciences LLC
122-25-067
EPA-accepted 08-24-98
Initial Printing

